

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

Doc Ref. **FP15**
Appl. No. 10/564,511
0 368 592
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89311472.8


(51) Int. Cl.⁵: **A01N 43/78, C07D 277/20,
C07D 277/60**

(22) Date of filing: 06.11.89

(30) Priority: 07.11.88 GB 8826035

(43) Date of publication of application:
16.05.90 Bulletin 90/20(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE(71) Applicant: **IMPERIAL CHEMICAL INDUSTRIES
PLC**
Imperial Chemical House, Millbank
London SW1P 3JF(GB)(72) Inventor: **Lewis, Terence**
18 Quintills Roman Hill
Bracknell Berkshire(GB)
Inventor: **Bansal, Harjinder Singh**
15 Chisbury Close
Bracknell Berkshire(GB)
Inventor: **Sunley, Raymond Leo**
1 Mallard Close Twyford
Berkshire(GB)Inventor: **Bartley, Michael Robert**
47 Blagrove Drive Wokingham
Berkshire(GB)Inventor: **Hepworth, Walter**
2 Dragons Hill Lyme Regis
Dorset(GB)Inventor: **Gilman, David John**
4 Gloucester Close Tytherington
Macclesfield Cheshire(GB)Inventor: **Kay, Ian Trevor**
'Chy-an-Ryn' Love Lane Mousehole
Penzance Cornwall(GB)Inventor: **Collins, David John**
10 Ardwell Close Crowthorne
Berkshire(GB)(74) Representative: **Downes, John Edward et al**
Imperial Chemical Industries PLC Legal
Department: Patents PO Box 6
Bessemer Road Welwyn Garden City
Hertfordshire AL7 1HD(GB)(54) **Herbicidal compositions.**

(57) A method of killing or controlling unwanted plants by applying to the plant on to a locus thereof an effective amount of a compound of formula (I)



or a salt thereof: wherein R^1 , R^2 , R^3 , R^4 and R^5 are independently selected from hydrogen, hydroxy, alkyl, alkoxy, alkylcarbonyl, halogen, nitrile, nitro, haloalkyl and haloalkoxy;
 R^6 is hydrogen, hydroxy, amino, lower alkyl, halogen, nitrile, haloalkyl, nitro, aryl, or a group $C(O)_m R^{10}$.

EP 0 368 592 A1

wherein R^{10} is hydrogen, alkyl, alkenyl, alkynyl or phenyl any of which may be optionally substituted and m is 1 or 2; R^7 is hydrogen, lower alkyl, haloalkyl, $C(O)_mR^{10}$ or halogen; or R^6 and R^7 together form an optionally substituted alkylene chain of 2 or 3 carbon atoms; R^8 is hydrogen, lower alkyl or halogen or the group R^7 and R^8 together form an oxo group; and the group CZR^9 is carboxy or an ester thereof, or R^9 is a group SR^{10} or $NR^{11}R^{12}$ wherein R^{11} is hydrogen or alkyl and R^{12} is hydrogen, alkyl, $S(O)_nR^{10}$ wherein n is 0, 1 or 2; or R^9 is a group $NR^{11}NR^{13}R^{14}$ wherein R^{11} , R^{13} and R^{14} are independently selected from hydrogen or alkyl; or R^9 is a group $- \overset{\oplus}{N}R^{11}NR^{13}R^{14}R^{20} X^{\ominus}$ wherein R^{11} , R^{13} , R^{14} and R^{20} are each hydrogen or alkyl, and x^- is an agriculturally acceptable anion, and Z is oxygen or sulphur.


Herbicidal compositions containing compounds of formula (I) are also claimed together with certain novel sub groups of compounds of formula (I) and processes for their preparation.

HERBICIDAL COMPOSITIONS

The present invention relates to the use of certain thiazole derivatives as herbicides and herbicidal compositions containing these compounds. Some of these compounds are novel and these, together with processes for their preparation form a further aspect of the invention.

A large number of thiazole derivatives are known in particular as pharmaceuticals (see for example British Patent Number 1,099,389). British Patent Number 1,022,750 and US Patent Number 3,418,331 show that some are also known as herbicides.

According to the present invention there is provided a method for killing or controlling unwanted plants by applying to the plants or to locus thereof a herbicidally effective amount of a compound of formula (I)



20 or a salt thereof: wherein R^1 , R^2 , R^3 , R^4 and R^5 are independently selected from hydrogen, hydroxy, alkyl, alkoxy, alkylcarbonyl, halogen, nitrile, nitro, haloalkyl and haloalkoxy;

R^6 is hydrogen, hydroxy, amino, mono- or di-alkyl amino, lower alkyl, halogen, nitrile, haloalkyl, nitro, aryl, or a group $C(O)_m R^{10}$ wherein R^{10} is hydrogen, alkyl, alkenyl, alkynyl or phenyl any of which may be optionally substituted and m is 1 or 2;

25 R^7 is hydrogen, lower alkyl, haloalkyl, $C(O)_m R^{10}$ or halogen; or R^5 and R^7 together form an optionally substituted alkylene chain of 2 or 3 carbon atoms;

R^8 is hydrogen, lower alkyl or halogen or the group R^7 and R^8 together form an oxo group; and the group CZR^9 is carboxy or an ester thereof, or R^9 is a group SR^{10} or $NR^{11}R^{12}$ wherein R^{11} is hydrogen or alkyl and R^{12} is hydrogen, optionally substituted alkyl, aryl, $S(O)_n R^{10}$ where n is 0, 1 or 2, or R^9 is a group $-NR^{11}NR^{13}R^{14}$ wherein R^{11} , R^{13} and R^{14} are independently selected from hydrogen or alkyl; or R^9 is a group $-NR^{11} \overset{+}{N}R^{13}R^{14}R^{20} X^-$ wherein R^{11} , R^{13} , R^{14} and R^{20} are each hydrogen or alkyl and X^- is an agriculturally acceptable anion; and Z is oxygen or sulphur.

Salts of the compounds of formula (I) are available when the compound includes a carboxy group. Suitable salts are agriculturally acceptable salts such as sodium, potassium, calcium, ammonium or sulphonium salts. Examples of ammonium salts are those of formula $NR^aR^bR^cR^d$ wherein R^a , R^b , R^c , and R^d , are independently selected from hydrogen or C_{1-10} alkyl optionally substituted with for example hydroxy.

Examples of sulphonium salts include those of formula $R^aR^bR^cS$ where R^a , R^b and R^c are independently selected from optionally substituted C_{1-10} alkyl.

40 As used herein the term "alkyl" includes straight or branched chains suitably containing from 1 to 10 carbon atoms, preferably from 1 to 6 carbon atoms. The expression "lower alkyl" refers to alkyl groups having from 1 to 3 carbon atoms. Similarly the terms "alkynyl" refer to unsaturated straight or branched chains having from 2 to 10, preferably from 2 to 6 carbon atoms.

45 In addition the term "alkoxy" refers to group O-alkyl as defined above. The terms "haloalkyl" and "haloalkoxy" refer to the described alkyl groups and alkoxy groups respectively which are substituted by at least one halogen atom such as fluorine, chlorine, bromine or iodine. A particular haloalkyl group is trifluoromethyl and a particular haloalkoxy group is trifluoromethoxy.

As used herein the term "aryl" includes phenyl and naphthyl.

50 Suitable groups R^1 , R^2 , R^3 , R^4 and R^5 are hydrogen, $C_{(1-6)}$ alkyl, $C_{(1-6)}$ alkoxy, lower alkyl carbonyl, halo $C_{(1-6)}$ alkyl, halo (C_{1-6}) alkoxy or halo wherein the halo groups are selected from fluorine, chlorine, bromine and iodine.

Particular examples of the groups R^1 , R^2 , R^3 , R^4 and R^5 are hydrogen, methyl, methoxy, trifluoromethyl trifluoromethoxy, fluoro, chloro, bromo, iodo or nitrile.

In a particular embodiment R^3 is other than hydrogen. Preferably R^3 is halogen such as chlorine and R^1 ,

R², R⁴ and R⁵ are hydrogen.

Suitably R⁶ is lower alkyl such as methyl or ethyl, halogen such as fluorine, chlorine, bromine or iodine, amino, hydroxy, nitro, nitrile, carboxy or lower alkyl esters thereof or trifluoromethyl. In particular R⁶ is bromine or chlorine.

Suitably R⁷ is selected from hydrogen, alkyl in particular lower alkyl, carboxy or lower alkyl esters thereof or haloalkyl such as trifluoromethyl or bromomethyl.

Suitably R⁸ is selected from hydrogen or methyl or haloalkyl such as trifluoromethyl.

Preferably both R⁷ and R⁸ are hydrogen.

When R⁶ and R⁷ form an optionally substituted alkylene chain, the substituents are suitably selected from those described above for R¹⁵ below.

When the group CZR⁹ is an ester of a carboxy group, it is suitably a group of formula CO₂R¹⁵ wherein R¹⁵ is alkyl, alkenyl, alkynyl or phenyl any of which may be optionally substituted.

Suitable optional substituents for R¹⁵ include one or more groups selected from halo such as fluoro, chloro, bromo or iodo; hydroxy; C₁₋₆ alkoxy; C₍₁₋₆₎ alkoxyC₍₁₋₆₎ alkoxy; aryl C₍₁₋₆₎ alkoxy; nitro; cycloalkyl; heterocyclic optionally substituted by oxo; nitrile; phenyl optionally substituted by nitro, halo such as chloro, alkoxy or carboxy or salts or C₁₋₆ alkyl esters thereof; or alkylsilyl groups such as trimethylsilyl.

As used herein the term "cycloalkyl" refers to cyclic alkyl groups containing from 3 to 10 ring carbon atoms. The term "heterocyclic" refers to rings of 3 to 10 atoms containing at least one atom selected from oxygen, nitrogen or sulphur.

When R¹⁵ is an alkenyl or alkynyl group, it suitably contains from 2 to 10 carbon atoms, preferably from 2 to 6 carbon atoms.

Preferred groups R⁹ are OR¹⁶ where R¹⁶ is hydrogen or unsubstituted C₁₋₆ alkyl.

Preferably Z is oxygen.

When R⁹ is a group SR¹⁰, R¹⁰ is preferably an alkyl group such as lower alkyl in particular ethyl. When R⁹ is a group NR¹¹R¹², R¹¹ and R¹², are suitably selected from hydrogen or lower alkyl such as methyl. R¹² may additionally be selected from phenyl, lower alkyl substituted by carboxy or lower alkyl esters thereof. R⁹ may also be a group -NR¹¹NR¹³R¹⁴ where R¹³ and R¹⁴ are independently hydrogen or lower alkyl such as methyl, or R⁹ may be a group -NR¹¹NR¹³R¹⁴R²⁰X[⊖].

Preferred groups NR¹¹R¹² are NH₂, N(CH₃)₂ or NHNH₂.

When R⁹ is a group -NR¹¹NR¹³R¹⁴R²⁰X[⊖], X[⊖] may be a halide anion, for example chloride, bromide or iodide.

Suitable substituents for R¹⁰ include those described above for R¹⁵.

Examples of compounds of formula (I) are set out in Tables I and II below. In all cases unless otherwise stated Z is oxygen.

TABLE I

COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
1	H	H	Br	H	H	H	H	H	NH ₂
2	H	H	CH ₃	H	H	H	H	H	OH
3	H	H	Cl	H	H	H	H	H	OH
4	H	H	Cl	H	H	Cl	H	H	OH
5	H	H	Br	H	H	H	H	H	OCH ₂ CH ₃
6	H	Cl	Cl	H	H	H	H	H	OH
7	H	H	Cl	H	H	H	H	H	OCH ₃
8	H	H	Cl	H	H	H	H	H	OCH ₂ CH ₃
9	H	H	Cl	H	H	Br	H	H	OH
10	H	CH ₃	Cl	H	H	H	H	H	OH
11	H	H	Cl	H	H	H	CH ₃	CH ₃	OH
12	H	OCH ₃	Cl	H	H	H	H	H	OH
13	H	H	Cl	H	H	H	H	H	NH-NH ₂
14	H	H	Cl	H	H	Cl	H	H	OCH ₃
15	H	H	Cl	H	H	Cl	H	H	O(CH ₂) ₂ CH ₃
16	H	H	Cl	H	H	Cl	H	H	NH ₂
17	H	H	Cl	H	H	Br	H	H	OCH ₃
18	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₂ CH ₃
19	H	H	Cl	H	H	Br	H	H	OCH ₂ CH ₃
20	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH ₃
21	H	H	Cl	H	H	Cl	H	H	NHNH ₂
22	H	H	H	H	H	H	H	H	N(CH ₃) ₂
23	H	H	Cl	H	H	H	H	H	N(CH ₃) ₂
24	H	H	H	H	H	H	CH ₃	H	N(CH ₃) ₂
25	H	H	Cl	H	H	H	H	H	NHCH ₂ CH ₃
26	H	H	H	H	H	H	H	H	N(CH ₂ CH ₃) ₂

TABLE I (Cont/D)

COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
27	H	H	H	H	H	H	H	H	NHCH ₂ CH ₃
28	H	H	H	H	H	Cl	H	H	N(CH ₃) ₂
29	Cl	H	H	H	H	H	H	H	CH ₂ CH ₃
30	Cl	H	H	H	H	H	H	H	N(CH ₃) ₂
31	H	H	H	H	H	H	CH ₃	CH ₃	N(CH ₃) ₂
32	Cl	H	H	H	Cl	H	H	H	N(CH ₃) ₂
33	Cl	H	H	H	Cl	H	CH ₃	H	N(CH ₃) ₂
34	Cl	H	H	H	H	H	CH ₃	CH ₃	N(CH ₃) ₂
35	CH ₃	H	H	H	H	H	H	H	N(CH ₃) ₂
36	CH ₃	H	H	H	H	H	CH ₃	H	N(CH ₃) ₂
37	Cl	H	H	H	H	H	CH ₃	H	N(CH ₃) ₂
38	CF ₃	H	H	H	H	H	H	H	OH
39	Cl	H	H	H	H	H	H	H	N(CH ₃) ₂
40	F	H	H	H	F	H	H	H	OCH ₂ CH ₃
41	F	H	H	H	F	H	H	H	OH
42	F	H	H	H	F	H	H	H	N(CH ₃) ₂
43	Cl	H	H	H	Cl	H	H	H	OCH ₂ CH ₃
44	Cl	H	H	H	F	H	H	H	OCH ₂ CH ₃
45	Br	H	H	H	H	H	H	H	OCH ₂ CH ₃
46	Br	H	H	H	H	H	H	H	OH
47	Br	H	H	H	H	H	H	H	N(CH ₃) ₂
48	Cl	H	H	H	F	H	H	H	OH
49	Cl	H	H	H	F	H	H	H	N(CH ₃) ₂
50	Cl	H	H	H	Cl	H	H	H	NHNH ₂
51	Cl	H	H	CF ₃	H	H	H	H	OCH ₂ CH ₃
52	Cl	H	H	CF ₃	H	H	H	H	OH
53	Cl	H	H	CF ₃	H	H	H	H	N(CH ₃) ₂

TABLE I (Cont/D)

5

10

15

20

25

30

35

40

45

50

55

COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
54	CH ₂ CH ₃	H	H	H	H	H	H	H	OCH ₂ CH ₃
55	H	Br	Br	H	H	H	H	H	OH
56	H	H	Cl	H	H	CH ₂ CH ₃	H	H	OCH ₂ CH ₃
57	H	H	Cl	H	H	CH ₃	H	H	OCH ₂ CH ₃
58	H	H	Cl	H	H	H	CH ₃	H	OCH ₂ CH ₃
59	H	H	Cl	H	H	H	CH(CH ₃) ₂	H	OCH ₂ CH ₃
60	H	H	Cl	H	H	Br	CH(CH ₃) ₂	H	OCH ₂ CH ₃
61	H	H	Cl	H	H	Cl	CH ₃	H	OCH ₂ CH ₃
62	H	H	Cl	H	H	Cl	H	H	N(CH ₃) ₂
63	H	H	Cl	H	H	Cl	H	H	NHN(CH ₃) ₂
64	H	H	Cl	H	H	Br	H	H	OCH(CH ₃) ₂
65	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₃ CH ₃
66	H	H	CF ₃	H	H	H	H	H	OCH ₂ CH ₃
67	H	H	I	H	H	H	H	H	OCH ₂ CH ₃
68	H	H	Cl	H	H	Cl	H	H	NHSO ₂ CH ₃
69	H	H	Cl	H	H	Br	H	H	NHC ₆ H ₅
70	H	H	I	H	H	Br	H	H	OCH ₂ CH ₃
71	H	H	Cl	H	H	Br	H	H	OCH ₂ CH ₂ OCH ₃
72	H	H	Cl	H	H	Br	H	H	OCH ₂ CH ₂ OH
73	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH ₂ CH(CH ₃) ₂
74	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH=CH ₂
75	H	H	Cl	H	H	Cl	H	H	OCH(CH ₂ CH ₃)CH ₂ CH ₃
76	H	H	Cl	H	H	Cl	H	H	OCH ₂ C≡CH
77	H	H	Cl	H	H	Cl	H	H	O(CH ₂) ₄ CH ₃
78	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH(CH ₃)- (CH ₂) ₃ CH(CH ₃) ₂

TABLE I (Cont/D)



COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
79	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH ₂ OCH ₂ (C ₆ H ₅)
80	H	H	Cl	H	H	Cl	H	H	OCH ₂ C(CH ₃) ₃
81	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH(CH ₃) ₂
82	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH(CH ₂ CH ₃)CH ₃
83	H	H	Cl	H	H	Cl	H	H	OCH(CH ₃)CH ₂ CH ₂ CH ₃
84	H	H	Cl	H	H	Cl	H	H	OCH(CH ₃)CH(CH ₃) ₂
85	H	H	Cl	H	H	Cl	H	H	OCH(CH ₃)CH ₂ CH ₃
86	H	H	Cl	H	H	Cl	H	H	OCH ₂ (C ₆ H ₁₁)
87	H	H	Cl	H	H	Cl	H	H	OCH(CH ₃)CH ₂ OCH ₃
88	H	H	Cl	H	H	Cl	H	H	O(CH ₂) ₅ CH ₃
89	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH ₂ CH(OCH ₃)CH ₃
90	H	H	Cl	H	H	Cl	H	H	OCH ₂ C ₆ H ₅
91	H	H	Cl	H	H	Cl	H	H	
92	H	H	Cl	H	H	Cl	H	H	O(CH ₂) ₇ CH ₃
93	H	H	Cl	H	H	Cl	H	H	
94	H	H	Cl	H	H	Cl	H	H	O(CH ₂) ₉ CH ₃
95	H	H	Cl	H	H	Cl	H	H	OC ₆ H ₅
96	H	H	Cl	H	H	Br	H	H	OCH ₂ C(CH ₃) ₃
97	H	H	Cl	H	H	Br	H	H	OCH(CH ₃)CH ₂ OCH ₃
98	H	H	Cl	H	H	Br	H	H	OC ₆ H ₅
99	H	H	Cl	H	H	Br	H	H	OCH ₂ C ₆ H ₅
x 100	H	H	Cl	H	H	Br	H	H	OCH ₂ C ₆ H ₁₁ Z

TABLE I (Cont/D)


COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
101	H	H	Cl	H	H	Br	H	H	
102	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₇ CH ₃
103	H	H	Cl	H	H	Cl	H	H	OC(CH ₃) ₃
104	H	H	Cl	H	H	Cl	H	H	OCH ₂ C(CH ₃) ₂ CH ₂ Cl
105	H	H	Cl	H	H	Cl	H	H	OCH ₂ Si(CH ₃) ₃
106	H	H	Cl	H	H	Cl	H	H	OC(CH ₃) ₂ CH ₂ Cl
107	H	H	Cl	H	H	Cl	H	H	OCH ₂ CH ₂ OCH ₂ CH ₂ OCH ₃
108	H	H	Cl	H	H	Cl	H	H	OCH ₂ CO ₂ CH ₂ CH ₃
109	H	H	Cl	H	H	Cl	H	H	OCH ₂ <u>CHOCH₂CH₂CH₂</u>
110	H	H	Cl	H	H	Cl	H	H	OCH(C ₆ H ₅)CO ₂ CH ₂ CH ₃
111	H	H	Cl	H	H	Br	H	H	<u>OCHCH₂CH₂OC=O</u>
112	H	H	Cl	H	H	Br	H	H	OCH(CH ₃)CO ₂ CH ₃
113	H	H	Cl	H	H	Br	H	H	OCH ₂ (C ₆ H ₅)CO ₂ CH ₂ CH ₃
114	H	H	Cl	H	H	Br	H	H	OCH ₂ CO ₂ CH ₂ CH ₃
115	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₂ O(CH ₂) ₂ OCH ₃
116	H	H	Cl	H	H	Br	H	H	OCH(C ₆ H ₅)CO ₂ CH ₃
117	H	H	Cl	H	H	Cl	H	H	OCH(C ₆ H ₅)CO ₂ CH ₃
118	H	H	Cl	H	H	Cl	H	H	OC(CH ₃) ₂ CF ₂ CF ₂ H

TABLE I (Cont/D)

COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
119	H	H	Cl	H	H	Cl	H	H	OCH(CH ₃)CO ₂ CH ₃
120	H	H	Cl	H	H	Cl	H	H	NHCH ₂ CO ₂ CH ₂ CH ₃
121	H	H	Cl	H	H	Cl	H	H	OCHCH ₂ CH ₂ OC=O
122	H	H	Cl	H	H	Cl	Br	H	OCH ₂ CH ₃
123	H	H	Cl	H	H	Cl	CO ₂ C ₆ H ₅	H	OCH ₂ CH ₃
124	H	H	Cl	H	H	Cl	Cl	H	OCH ₂ CH ₃
125	H	H	Cl	H	H	Cl	= O	H	OCH ₂ CH ₃
126	H	H	Cl	H	H	NO ₂	H	H	OCH ₃
127	H	H	F	H	H	Br	H	H	OCH ₃
128	H	H	F	H	H	Br	H	H	OCH ₂ CH ₃
129	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₄ CH ₃
130	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₅ CH ₃
131	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₇ CH ₃
132	H	H	Cl	H	H	Cl	F	H	OCH ₂ CH ₃
133*	H	H	Cl	H	H	Br	H	H	OCH ₃
134	F	H	H	H	F	Br	Br	H	OCH ₂ CH ₃
135	H	CH ₃	H	H	H	H	H	H	OCH ₃
136	H	H	F	H	H	H	H	H	OCH ₃
137	H	H	F	H	H	NO ₂	H	H	OCH ₃
138	H	H	Cl	H	H	Br	H	H	O(CH ₂) ₂ CH ₃
139	H	H	F	H	H	H	H	H	OCH ₃
140	H	Cl	H	H	H	Br	H	H	OH
141	H	H	H	H	H	Br	H	H	OCH ₂ CH ₃
142	H	H	H	H	H	Br	H	H	OCH ₃
143	H	Cl	H	H	H	Br	H	H	OCH ₂ CH ₃

* 2 = sulphur


TABLE I (Cont/D)

COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
144	H	Cl	H	H	H	Br	H	H	OCH ₃
145	H	H	Cl	H	H	Br	Br	H	OCH ₃
146	H	H	Cl	H	H	Br	Br	H	OCH ₂ CH ₃
147	H	H	Br	H	H	Br	H	H	OCH ₃
148	H	H	Br	H	H	Br	H	H	OCH ₂ CH ₃
149	H	H	Br	H	H	Br	H	H	O(CH ₂) ₂ CH ₃
150	H	H	Br	H	H	Br	H	H	OH
151	H	H	CF ₃	H	H	Br	H	H	OH
152	H	H	CF ₃	H	H	Br	H	H	OCH ₃
153	H	H	CF ₃	H	H	Br	H	H	OCH ₂ CH ₃
154	H	H	Cl	H	H	NH ₂	H	H	OCH ₃
155	F	H	F	H	H	Br	H	H	OH
156	H	F	F	H	H	Br	H	H	OH
157	F	H	F	H	H	Br	H	H	OCH ₃
158	F	H	F	H	H	Br	H	H	OCH ₂ CH ₃
159	H	F	F	H	H	Br	H	H	OCH ₃
160	H	F	F	H	H	Br	H	H	OCH ₂ CH ₃
161	H	H	NO ₂	H	H	H	H	H	OH
162	H	H	H	H	H	Br	H	H	OH
163	H	H	F	H	H	Br	H	H	OH
164	F	H	H	H	F	Br	H	H	N(CH ₃) ₂
165	H	H	Cl	H	H	Br	H	H	NHN(CH ₃) ₂
166	H	H	Cl	H	H	Br	H	H	\oplus NHN(CH ₃) ₃ I [⊖]
167	H	H	CH ₃	H	H	Br	H	H	OCH ₃
168	H	H	Cl	H	H	Br	H	H	SCH ₂ CH ₃

TABLE I (Cont/D)


COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹
169	H	H	OCH ₃	H	H	Br	H	H	OCH ₃
170	H	H	OCH ₃	H	H	Br	H	H	OH
171	H	H	OCF ₃	H	H	Br	H	H	OCH ₃
172	H	H	COCH ₃	H	H	Br	H	H	OH
173	H	H	Cl	H	H	H	F	H	OCH ₃
174	H	H	OH	H	H	Br	H	H	OCH ₃
175	H	H	Cl	H	H	CO ₂ CH ₃	H	H	OCH ₃
176	H	H	Cl	H	H	CO ₂ H	H	H	OH
177	H	H	Cl	H	H	OH	H	H	OH
178	H	H	Cl	H	H	CH ₃	CH ₃	H	OCH ₃
179	H	H	Cl	H	H	CH ₃	CH ₂ Br	H	OCH ₃

TABLE II



COMPOUND NO.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁹	A
180	H	H	Cl	H	H	H	OCH ₂ CH ₃	-CH ₂ -
181	H	H	Cl	H	H	H	OH	-CH ₂ -
182	H	H	Cl	H	H	H	OCH(CH ₃) ₂	-CH ₂ -
183	H	H	H	H	H	H	OCH ₂ CH ₃	-CH ₂ -
184	H	H	C(CH ₃) ₃	H	H	H	OCH ₂ CH ₃	-CH ₂ -
185	H	H	CF ₃	H	H	H	OCH ₂ CH ₃	-CH ₂ -
186	H	H	Cl	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
187	H	H	H	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
188	H	Cl	F	H	H	H	OCH ₂ CH ₃	-CH ₂ -
189	H	Cl	F	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
190	H	H	Br	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
191	Cl	H	H	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
192	H	H	Br	H	H	H	OCH ₂ CH ₃	-CH ₂ -
193	Cl	H	H	H	Cl	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
194	H	H	CH ₃	H	H	H	OCH ₂	-CH ₂ -CH ₂ -
195	Cl	H	Cl	H	H	H	OCH ₂ CH ₃	-CH ₂ -CH ₂ -
196	Cl	H	H	H	F	H	OCH ₂ CH ₃	-CH ₂ -
197	H	H	F	H	H	H	OCH ₂ CH ₃	-CH ₂ -


Many of the compounds of formula (I) are known compounds and can be prepared by known methods for example as set out in GB Patent Number 1,099,389. An example of preparation is set out in Scheme A below. The reaction conditions employed in each step of the process are conventional.

Scheme A

In scheme A, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 and Z are as defined in relation to formula (I), Y is a leaving group such as chloro, and R^6 is H, lower alkyl, or $-\text{CO}_2\text{R}^{10}$.

Certain of the compounds of formula (I) are novel and these form a further aspect of the invention.


In particular according to the invention there is provided a compound of formula (II)



15 wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , R^9 and Z are as defined in relation to formula (I) and R^{17} is halogen, nitro, hydroxy, amino, mono- or di-alkylamino or nitrile.

In a preferred embodiment R^{17} is halogen such as fluoro, chloro, bromo, or iodine preferably bromo.

Compounds of formula (II) where R^{17} is halogen can be prepared by reacting a compound of formula (III)



35 wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 and Z are as defined in relation to formula (I); with chlorine or bromine and thereafter if necessary converting the group $CZOH$ to a different group CZR^9 as defined in relation to formula (I).

The reaction is suitably carried out in a solvent such as glacial acetic acid and/ or chloroform at temperatures of from 20°-80°C.

40 Compounds of formula II where R^{17} is nitrile can be prepared by subjecting the products of Scheme A wherein R^6 is $-CO_2R^{10}$ to conventional procedures for transforming an alkoxy-carbonyl group to a $-CN$ group.

Compounds of formula (II) where R^{17} is nitro can be prepared by reacting an ester derivative of a compound of formula (III) with a mild nitrating agent such as nitronium tetrafluoroborate. The reaction is suitably effected in an organic solvent such as CH_3CN . Low temperatures of from 0°C - 15°C are suitably employed.


45 These nitro compounds can be converted to compounds of formula (II) where R^{17} is amino by hydrogenation using conventional conditions. For instance, the compound can be reacted with hydrogen in the presence of a catalyst such as palladium on charcoal. The reaction is suitably effected in an organic solvent such as methanol or dichloromethane. Alkylation of the amino compound for example by reaction with an alkyl halide in the presence of base gives a compound of formula (II) where R^{17} is the mono- or di-alkyl amino group.

Compounds of formula (III) are known compounds and fall within the definition of formula (I), or can be prepared from known compounds by conventional methods.

55 Conversion of the group $CZOH$ to a CZR^9 group can be carried out by conventional methods. One simple method for preparing a compound of formula (I) where R^9 is OR^{10} and R^{10} is lower alkyl is by refluxing the compound in an appropriate alcohol of formula $R^{19}OH$ wherein R^{19} is alkyl such as methanol or ethanol, in the presence of a catalytic amount of acid such as sulphuric acid.


Alternatively the compound can be reacted with an appropriate alcohol, dicyclohexylcarbodiimide (DCC)- as a coupling agent, in the presence of a base such as DMAP. The reaction may be carried out in an organic solvent such as dichloromethane or chloroform. Low temperatures, for example from -5 to 20°C are suitably employed.

Other compounds of formula (I) which are novel and which form an additional aspect of the invention are compounds of formula (V)




wherein R¹, R², R³, R⁴, R⁵, R⁸, R⁹ and Z are as defined in relation to formula (I)

These compounds are prepared by reacting a compound of formula (VI)



wherein R¹, R², R³, R⁴ and R⁵ are as defined in relation to formula (I); with a compound of formula (VII)



wherein R⁸, R⁹ and Z are as defined in relation to formula (I), and X¹ is a leaving group such as chlorine or bromine.

The reaction is suitably carried out in an organic solvent such as ethanol or toluene at elevated temperatures from 60 to 120°C

Compounds of formula (VI) and (VII) are either known compounds or they can be prepared from known compounds by conventional methods.

The compounds of formula (I) are active as herbicides and therefore, in a further aspect the invention provides a process for severely damaging or killing unwanted plants which process comprises applying to the plants, or to the growth medium of the plants, an effective amount of a compound of formulae (I) as hereinbefore defined.

The compounds of formula (I) are active against a broad range of weed species including monocotyledonous and dicotyledonous species. The compounds may show a useful selectivity into monocotyledonous crops, in particular rice.

The compounds of formula (I) may be applied directly to the plant (post-emergence application) or to the soil before the emergence of the plant (pre-emergence application).

The compounds of formulae (I) may be used on their own to inhibit the growth of, severely damage, or kill plants but are preferably used in the form of a composition of the invention.

Compositions containing compounds of formula (I) include both dilute compositions, which are ready for immediate use, and concentrated compositions, which require to be diluted before use, usually with water.

- 5 Preferably the compositions contain from 0.01% to 90% by weight of the active ingredient. Dilute compositions ready for use preferably contain from 0.01% to 2% of active ingredient, while concentrated compositions may contain from 20 to 90% of active ingredient, although from 20 to 70% is usually preferred.

- 10 The solid compositions may be in the form of granules, or dusting powders wherein the active ingredient is mixed with a finely divided solid diluent, eg, kaolin, bentonite, kieselguhr, dolomite, calcium carbonate, talc, powdered magnesia Fuller's earth and gypsum. They may also be in the form of dispersible powders or grains, comprising a wetting agent to facilitate the dispersion of the powder or grain in liquid. Solid compositions in the form of a powder may be applied as foliar dusts.

- 15 Liquid compositions may comprise a solution or dispersion of an active ingredient in water optionally containing a surface-active agent, or may comprise a solution or dispersion of an active ingredient in a water-immiscible organic solvent which is dispersed as droplets in water.

- Surface-active agents may be of the cationic, anionic, or non-ionic type. The cationic agents are, for example, quaternary ammonium compounds (eg cetyltrimethylammonium bromide). Suitable anionic agents are soaps; salts or aliphatic mono esters of sulphuric acid, for example sodium lauryl sulphate; and salts of 20 sulphonated aromatic compounds, for example sodium dodecylbenzenesulphonate, sodium, calcium and ammonium lignosulphonate, butylnaphthalene sulphonate, and a mixture of the sodium of diisopropyl and triisopropynaphthalenesulphonic acid. Suitable non-ionic agents are the condensation products of ethylene oxide with fatty alcohols such as oleyl alcohol and cetyl alcohol, or with alkylphenols such as octyl- or nonyl-phenol or octyl-cresol. Other non-ionic agents are the partial esters derived from long chain fatty 25 acids and hexitol anhydrides, for example sorbitan monolaurate; the condensation products of the partial ester with ethylene oxide; and the lecithins.

- The aqueous solutions or dispersions may be prepared by dissolving the active ingredient in water or an organic solvent optionally containing wetting or dispersing agent(s) and then, when organic solvents are used, adding the mixture so obtained to water optionally containing wetting or dispersing agent(s). Suitable 30 organic solvents include, for example, ethylene di-chloride, isopropyl alcohol, propylene glycol, diacetone alcohol, toluene, kerosene, methylnaphthalene, the xylenes and trichloroethylene.

- The compositions for use in the form of aqueous solutions or dispersions are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, and the concentrate is then 35 diluted with water before use. The concentrates are usually required to withstand storage for prolonged periods and after such storage, to be capable of dilution with water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Concentrates conveniently contain 20-90%, preferably 20-70%, by weight of the active ingredient(s). Dilute preparations ready for use may contain varying amounts of the active ingredient(s) depending upon the intended purpose; amounts of 0.01% to 10.0% and preferably 0.1% to 2%, by weight of active ingredient(s) 40 are normally used.

- A preferred form of concentrated composition comprising the active ingredient which has been finely divided and which has been dispersed in water in the presence of a surface-active agent and a suspending agent. Suitable suspending agents are hydrophilic colloids and include, for example, polyvinylpyrrolidone and sodium carboxymethylcellulose, and the vegetable gums, for example gum acacia and gum tragacanth. 45 Preferred suspending agents are those which impart thixotropic properties too, and increase the viscosity of the concentrate. Examples of preferred suspending agents include hydrated colloidal mineral silicates, such as montmorillonite, beidellite, nontronite, hectorite, saponite, and saucorite. Bentonite is especially preferred. Other suspending agents include cellulose derivatives and polyvinyl alcohol.

- The rate of application of the compounds of the invention will depend on a number of factors including, 50 for example, the compound chosen for use, the identity of the plants whose growth is to be inhibited, the formulations selected for use and whether the compound is to be applied for foliage or root uptake. As a general guide, however, an application rate of from 0.01 to 10 kilogrammes per hectare is suitable.

- The compositions of the invention may comprise, in addition to one or more compounds of formula (I) or (II) one or more compounds not of the invention but which possess biological activity. Accordingly in yet 55 a still further embodiment the invention provides a herbicidal composition comprising a mixture of at least one herbicidal compound of formula (I) as hereinbefore defined with at least one other herbicide.

The other herbicide may be any herbicide not having the formula (I). It will generally be a herbicide having a complementary action in the particular application.

For example it may be desirable in certain circumstances to use the compound of formula (I) in admixture with a contact herbicide.

Examples of useful complementary herbicides include:

A. benzo-2,1,3-thiadiazin-4-one-2,2-dioxides such as 3-isopropylbenzo-2,1,3-thiadiazin-4-one-2, 2-dioxide (bentazon);

B. hormone herbicides, particularly the phenoxy alkanolic acids such as 4-chloro-2-methylphenoxy acetic acid (MCPA), S-ethyl 4-chloro-O-tolyloxy thio-acetate (MCPA-thioethyl), 2-(2,4-dichlorophenoxy) propionic acid (dichlorprop), 2,4,5-trichlorophenoxyacetic acid (2,4,5-T), 4-(4-chloro-2-methylphenoxy)butyric acid (MCPB), 2,4-dichlorophenoxyacetic acid (2,4-D), 4-(2,4-dichlorophenoxy)butyric acid (2,4-DB), 2-(4-chloro-2-methylphenoxy) propionic acid (mecoprop), 3,5,6-trichloro-2-pyridyloxyacetic acid (trichlopyr), 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid (fluroxypyr), 3,6-dichloropyridine-2-carboxylic acid (clopyralid), and their derivatives (eg. salts, esters and amides);

C. 1,3 dimethylpyrazole derivatives such as 2-[4-(2,4-dichlorobenzoyl) 1,3-dimethylpyrazol-5-yloxy] acetophenone (pyrazoxyfen), 4-(2,4-dichlorobenzoyl)1,3-dimethylpyrazol-5-yltoluene suphonate (pyrazolate) and 2-[4-(2,4-dichloro-m-toluoxy)-1,3-dimethylpyrazol-5-yloxy]-4-methylacetophenone (benzofenap);

D. Dinitrophenols and their derivatives (eg. acetates such as 2-methyl-4,6-dinitrophenol (DNOC), 2-*t*-butyl-4,6-dinitrophenol (dinoterb), 2-secbutyl-4,6-dinitrophenol (dinoseb) and its ester, dinoseb acetate;

E. dinitroaniline herbicides such as N',N'-diethyl-2,6-dinitro-4-trifluoromethyl-m-phenylenediamine (dinitramine), 2,6-dinitro-N,N-dipropyl-4-trifluoro-methylaniline (trifluralin),

N-ethyl-N-(2-methylallyl)-2,6-dinitro-4-trifluoromethylaniline (ethalflurolin), N-(1-ethylpropyl)-2,6-dinitro-3,4-xylidine (pendimethalin); and 3,5-dinitro-N⁴, N⁴-dipropylsulphanilamide (oryzalin);

F. arylurea herbicides such as N'-(3,4-dichlorophenyl)-N,N-dimethylurea (diuron), N,N-dimethyl-N'-[3-(trifluoromethyl) phenyl]urea (flumeturon), 3-(3-chloro-4-methoxyphenyl)-1, 1-dimethylurea (metoxuron), 1-butyl-3-(3,4-dichlorophenyl)-1-methylurea (neburon), 3-(4-isopropylphenyl)-1,1-dimethylurea (isoproturon), 3-(3-chloro-p-tolyl)-1,1-dimethylurea (chlorotoluron), 3-[4-(4-chlorophenoxy, phenyl)-1, 1-dimethylurea (chloroxuron), 3-(3,4-dichlorophenyl)-1-methylurea (linuron), 3-(4-chlorophenyl)-1-methoxy-1-methylurea (monolinuron), 3-(4-bromo-4-chlorophenyl)-1-methoxy-1-methylurea (chlorobromuron), 1-(1-methyl-1-phenylethyl)-3-p-tolylurea (daimuron), and 1-benzothiazol-2-yl-1,3-dimethylurea (methabenzthiazuron);

G. phenylcarbamoyloxyphenylcarbamates such as 3-[methoxycarbonylamino]phenyl (3-methyl-phenyl)-carbamate (phenmedipham) and 3-[ethoxycarbonylamino]-phenyl phenylcarbamate (desmedipham);

H. 2-phenylpyridazin-3-ones such as 5-amino-4-chloro-2-phenylpyridazin-3-one (pyrazon), and 4-chloro-5-methylamino-2-(, -trifluoro-m-tolyl) pyridazin-3(2H)-one (norflurazon);

I. uracil herbicides such as 3-cyclohexyl-5,6-trimethyleneuracil (lenacil), 5-bromo-3-sec-butyl-6-methyl-uracil (bromacil) and 3-*t*-butyl-5-chloro-6-methyl-uracil (terbacil);

J. triazine herbicides such as 2-chloro-4-ethylamino-6-(*i*-propylamino)-1,3,5-triazine (atrazine), 2-chloro-4,6-di(ethylamino)-1,3,5-triazine (simazine), 2-azido-4-(*i*-propylamino)-6-methylthio-1,3,5-triazine (aziprotryne), 2-(4-chloro-6-ethylamino-1,3,5-triazin-2-ylamino)-2-methylpropionitrile (cyanazine), N², N⁴-diisopropyl-6-methylthio-1,3,5-triazine-2,4-diamine (prometryne), N²-(1,2-dimethylpropyl)-N⁴-ethyl-6-methylthio-1,3,5-triazine-2,4-diamine (dimethametryne), N²,N⁴-diethyl-6-methylthio-1,3,5-triazine-2,4-diamine (simetryne), and N²-*tert*-butyl-N⁴-ethyl-6-methylthio-1,3,5-triazine-2,4-diamine (terbutryne);

K. phosphorothioate herbicides such as S-2-methylpiperidinocarbonyl-methyl O,O-dipropyl phosphorodithioate (piperophos), S-2-benzenesulphonamidoethyl O,O-di isopropyl phosphonodithioate (bensulide), and O-ethyl O-6-nitro-m-tolyl sec-butylphosphoamidothioate (butamifos);

L. thiolcarbamate herbicides such as S-ethyl N-cyclohexyl-N-ethyl(thiocarbamate) (cycloate), S-propyl dipropyl-thiocarbamate (vernolate), S-ethyl-azepine-1-carbothioate (molinate), S-4-chlorobenzyl diethyl-thiocarbamate (thiobencarb), S-ethyl diisobutyl-thiocarbamate (butylate) *, S-ethyl diisopropylthiocarbamate (EPTC) *, S-2,3,3-trichloroallyl di-isopropyl (thiocarbamate) (tri-allate), S-2, 3-dichloroallyl di-isopropyl (thiocarbamate) (diallate), S-benzyl 1,2-dimethylpropyl (ethyl) thiocarbamate (esprocarb), S-benzyl di(sec-butyl)-thiocarbamate (triocarbamil), 6-chloro-3-phenylpyridazin 4-yl S-octyl thiocarbamate (pyridate), and S-1-methyl-1-phenylethylpiperidine -1-carbothioate (dimepiperate);

M. 1,2,4-triazin-5-one herbicides such as 4-amino-4,5-dihydro-3-methyl-6-phenyl-1,2,4-triazine-5-one (metamitron) and 4-amino-6-*t*-butyl-4,5-dihydro-3-methylthio-1,3,4-triazin-5-one (metribuzin);

N. benzoic acid herbicides such as 2,3,6-trichlorobenzoic acid (2,3,6-TBA), 3,6-dichloro-2-methoxybenzoic acid (dicamba) and 3-amino-2, 5-dichloro benzoic acid (chloramben);

O. anilide herbicides such as 2-chloro-2',6'-diethyl-N-(2-propoxyethyl)acetanilide (pretilachlor), N-

* These compounds are preferably employed in combination with a safener such as 2,2-dichloro-N,N-di-2-propenylacetamide (dichlormid)

- butoxymethyl-chloro-2', 6'-diethylacetanilide (butachlor), the corresponding N-methoxy compound (alachlor), the corresponding N-i-propyl compound (propachlor), 3',4'-dichloropropionilide (propanil), 2-chloro-N-[pyrazol-1-ylmethyl]acet-2'-6'-xylidide (metazachlor), 2-chloro-6'-ethyl-N-(2-methoxy-1-methylethyl)-acet-0-toluidide (metolachlor), 2-chloro-N-ethoxymethyl-6'-ethylacet-0-toluidide (acetochlor), and 2-chloro-N-(2-methoxyethyl)acet-2',6'-xylidide (dimethachlor);
- 5 P. dihalobenzonitrile herbicides such as 2,6-dichloro-benzonitrile (dichlobenil), 3,5-dibromo-4-hydroxy-benzonitrile (bromoxynil) and 3,5-diiodo-4-hydroxy-benzonitrile (ioxynil);
- Q. haloalkanoic herbicides such as 2,2-dichloropropionic acid (dalapon), trichloroacetic acid (TCA) and salts thereof;
- 10 R. diphenylether herbicides such as ethyl 2-[5-(2-chloro-trifluoro-p-tolyloxy)-2-nitrobenzoyloxy] propionate (lactofen), D-[5-(2-chloro- , , -trifluoro-p-tolyloxy)-2-nitrobenzoyl] glycolic acid (fluroglycofen) or salts or ester thereof, 2,4-dichlorophenyl-4-nitrophenyl ether (nitrofen), methyl-(2,4-dichlorophenoxy)-2-nitrobenzoate (bifenox), 2-nitro-5-(2-chloro-4-trifluoromethyl-phenoxy) benzoic acid (acifluorfen) and salts and esters thereof, 2-chloro-4-trifluoromethylphenyl 3-ethoxy- 4-nitrophenyl ether (oxyfluorfen) and 5-(2-chloro-4-(trifluoromethyl)phenoxy)-N-(methylsulfonyl)-2-nitrobenzamide (fomesafen); 2,4,6-trichlorophenyl 4-nitrophenyl ether (chlornitrofen) and 5-(2,4-dichlorophenoxy)-2-nitroanisole (chlomethoxyfen);
- 15 S. phenoxyphenoxypropionate herbicides such as (RS)-2-[4-(2,4-dichloro-phenoxy)phenoxy] propionic acid (diclofop) and esters thereof such as the methyl ester, 2-[4-(5-trifluoromethyl)-2-(pyridinyl)oxy] phenoxypropanoic acid (fluazifop) and esters thereof, 2-[4-(3-chloro-5-trifluoro-methyl)-2-pyridinyl]oxy)phenoxypropanoic acid (haloxyfop) and esters thereof, 2-[4-(6-chloro-2-quinoxalinyloxy)phenoxy]propanoic acid (quizalofop) and esters thereof and (±)-2-[4-(6-chlorobenzoxazol-2-yloxy)phenoxy]propionic acid (fenoxaprop) and esters thereof such as the ethyl ester;
- 20 T. cyclohexanedione herbicides such as 2,2-dimethyl-4,6-dioxo-5-(1-((2-propenyloxy)imino)butyl) cyclohexane carboxylic acid (alloxydim) and salts thereof, 2-(1-ethoxyimino) butyl-5-(2-(ethylthio)propyl)-3-hydroxy-2-cyclohexan-1-one(ethoxydim), 2-(1-ethoxyimino) butyl-3 -hydroxy-5-thian-3-ylcyclohex-2-enone (cycloxydim) 2-[1-(ethoxyimino)propyl]-3-hydroxy-5-mesitylcyclohex-2-enone(tralkoxydim), and (±) -2- (E)-1-[(E)-3-chloroallyloximino]propyl -5-[2-(ethylthio)propyl]-3-hydroxy-cyclohex-2-enone (clethodim);
- U. sulfonyl urea herbicides such as 2-chloro-N (4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl benzenesulphonamide (chlorosulfuron), methyl 2-(((4,6-dimethyl-2-pyrimidinyl)amino)carbonyl)amino)-sulphonylbenzoic acid (sulfometuron), 2-(((3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)carbonyl) amino)-sulphonyl)benzoic acid (metsulfuron) and esters thereof; -(4,6-dimethoxypyrimidin-2-ylcarbamoylsulphamoyl)-O-toluic acid (benzsulfuron) and esters thereof such as the methyl 3-[3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-ureidosulphonyl]thiophene-2-carboxylate (DPX-M6313), 2-(4-chloro-6-methoxy pyrimidin-2-yl) carbamoylsulphamoyl benzoic acid (chlorimuron) and esters such as the ethyl ester thereof 2-(4,6-dimethoxypyrimidin-2-ylcarbamoylsulphamoyl)-N, N-dimethylnicotinamide, 2-[4,6-bis(difluoromethoxy) pyrimidin-2-ylcarbamoylsulphamoyl] benzoic acid (pirimisulfuron) and esters such as the methyl ester thereof, 2-[3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-3-methylureidosulphonyl] benzoic acid esters such as the methyl ester thereof (DPX-LS300) and 5-(4,6-dimethoxypyrimidin-2-ylcarbamoylsulphamoyl)-1-methylpyrazole-4-carboxylic acid (pyrazosulfuron);
- 30 V. imidazolidinone herbicides such as 2-(4,5-dihydro-4-isopropyl-4-methyl-5-oxoimidazol-2-yl) quinoline-3-carboxylic acid (imazaquin), methyl 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluate and p-toluate isomer(imazamethabenz), 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl) nicotinic acid (imazapyr) and isopropylammonium salts thereof, (RS)-5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid (imazethapyr);
- 35 W. arylanilide herbicides such as benzoyl-N-(3-chloro-4-fluorophenyl)-L-alanine (flamprop) and esters thereof, ethyl N-benzoyl-N-(3,4-dichlorophenyl)-DL-alaninate (benzoylprop-ethyl), N-(2,4-difluorophenyl)-2-(3-trifluoromethyl)phenoxy)-3-pyridinecarboxamide (diflufenican); and
- X. amino acid herbicides such as N-(phosphonomethyl)-glycine (glyphosate) and DL-homoalanin-4-yl-(methyl)-phosphinic acid (phosphinothricin) and their salts and esters, trimethylsulfonium N-(phosphonomethyl)-glycine (sulphosate), and bilanafos;
- 40 Y. organoarsenical herbicides such as monosodium methanearsonate (MSMA);
- Z. herbicidal amide derivative such as (RS)-N,N-diethyl-2-(1-naphthylloxypropionamide) (napropamide), 3,5-dichloro-N-(1,1-dimethylpropynyl)benzamide (propyzamide), (R)-1-(ethylcarbamoyl)ethyl carbanilate (carbetamide), N-benzyl-N- isopropylpivalamide (tebutam), (RS)-2-bromo- N-(, -dimethylbutylamide (bromobutide), N-[3-(1-ethyl-1-methylpropyl)-isoxazol-5-yl] 2,6-dimethoxybenzamide, (isoxaben), N-phenyl-2-(2-naphthylloxy) propionamide (naproanilide), N,N-dimethyl-diphenylacetamide (diphenamid), and N-(1-naphthyl)-phthalamic acid (naptalam);
- 45 AA. miscellaneous herbicides including 2-ethoxy-2,3-dihydro-3, 3-dimethylbenzofuran methanesul-

fonate (ethofumesate), 7-oxabicyclo (2.2.1)heptane,1-methyl-4-(1-methylethyl)-2-(2-methylphenylmethoxy)-exo (cinmethylin), 1,2-dimethyl-3,5-diphenylpyrazolium ion (difenzoquat) and salts thereof such as the methyl sulphate salt, 2-(2-chlorobenzyl)-4,4-dimethyl-1,2-oxazolidin -3-one (clomazone), 5-tert-butyl-3-(2,4-dichloro-5-isopropoxyphenyl)-1,3,4-oxadiazol-2(3H)-one (oxadiazon), 3,5-dibromo-4-hydroxy benzaldehyde
 5 2,4-dinitrophenyloxime (bromofenoxim), 4-chlorobut-2-ynyl-3-chlorocarbaniolate (barban), (RS)-2-(3,5-dichlorophenyl)-2-(2,2, 2-trichloroethyl)oxirane (tridiphane), (3RS,4RS; 3RS,4SR)-3-chloro-4-chloromethyl-1-(, , -trifluoro-m-totyl)-2-pyrrolidone (in the ratio 3:1) (fluorodichloridone), dichloroquinoline 8-carboxylic acid (quinchlorac) and 2-(1,3-benzothiazol-2-yl- oxy)-N-methylacetanilide (mefanacet);

BB. Examples of useful contact herbicides include:

10 bipyridylum herbicides such as thos in which the active entity is the 1,1'-dimethyl-4,4'-dipyridylum ion (paraquat) and those in whcih the active entity is the 1,1'-ethylene-2,2'-dipyridylum ion (diquat); The following examples are given by way of illustration only.

EXAMPLE 1

Preparation of 5-Bromo-2-(4-chlorophenyl) thiazol-4-yl-acetic acid (Compound No. 9 in Table 1).

20 Bromine (15.9g, 5.15ml, 0.1mol) was added dropwise to 2-(4-chlorophenyl) thiazol-4-yl acetic acid (24g, 0.096mol) suspended in glacial acetic acid (200ml) and the resulting mixture was heated at 60-70°C for 5 hours. The reaction mixture was cooled to 0°C (ice bath) and the solid which separated was filtered and washed with CHCl₃. The residue was then washed with cold ethanol to afford the desired product (24g,
 25 75%)

¹H NMR (DMSO-d₆) 3.75 (2H, S, CH₂CO) and 7.7 (4H, AB quartet, aromatic C-H)

EXAMPLE 2

Preparation of Methyl-5-bromo-2-(4-chlorophenyl thiazol-4-yl acetic ester (Compound No. 17 in Table 1).

35 The reaction as substantially described in Example 1 was carried out up to the cooling step. The solid which separated was filtered and washed with cold acetic acid. The crude 5-bromo-2-(4-chlorophenyl) thiazol-4-yl acetic acid without further purification was dissolved in hot methanol (100ml) and the resulting reaction mixture heated under reflux for 20 mins (the esterification was monitored by silica TLC, eluting in CH₂Cl₂). The reaction mixture was allowed to cool to room temperature, and the solid which separated out
 40 of solution was filtered and air dried to yield the desired product.

¹H NMR (CDCl₃) 3.65 (3H, s, OCH₃), 3.8 (2H, s, CH₂CO) and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 3

Preparation of Ethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester (Compound No. 19 in Table 1).

50 5-Bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid, (prepared as described in Example 1 but without recrystallisation from ethanol) was dissolved in a mixture of hot ethanol (100ml) and conc. H₂SO₄ (4 drops, 0.1ml) and the resulting solution was heated under reflux for 40 min. The reaction mixture was allowed to cool to room temperature and the solid which separated out of solution was filtered, and washed with cold ethanol to yield the desired product (24.2g,71%) m.p. 75-77 °C,

55 ¹H NMR (CDCl₃) 1.3 (3H,t,CH₂CH₃) 3.9 (2H,s,CH₂CO) 4.25 (2H,q,CH₂CH₃), and 7.55 (4H,AB, quartet, aromatic C-H).

EXAMPLE 4

5 Preparation of n-Propyl-5-bromo-2-(4-chlorophenyl) thiazol-4-yl acetic ester (Compound No. 18 in Table 1).

5-Bromo-2-(4-chlorophenyl) thiazol-4-yl acetic acid (prepared as described in Example 1 but without recrystallisation from ethanol) was dissolved in a mixture of hot n-propanol (100ml) and concentrated sulphuric acid (4 drops, 0.1ml) and the resulting solution was heated under reflux for 6 hours. The reaction mixture was cooled to room temperature and poured into H₂O (25ml). The solution was neutralised (NaHCO₃), and the precipitate filtered to yield the desired product (1.33g, 60%)
¹H NMR (CDCl₃) 0.95 (3H, t, CH₂CH₂CH₂CH₃), 1.35 (2H, hexet, CH₂CH₂CH₂CH₃), 1.6 (2H, quintet, CH₂CH₂CH₂CH₃), 3.9 (2H, s, CH₂CO), 4.15 (2H, t, OCH₂CH₂CH₂CH₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 6

20 Preparation of 5-Chloro-2-(4-chlorophenyl) thiazol-4-yl acetic ester (Compound No. 4 in Table I).

Chlorine gas was bubbled through a stirred suspension of 2-(4-chlorophenyl) thiazol-4-yl acetic acid (10g, 39mmol) in glacial acetic acid (150ml). After 1min the thiozolyacetic acid had dissolved and after 2 min of bubbling a precipitate began to form. Chlorine was bubbled for a total of 20min, and the reaction mixture allowed to stir for a further 30 minutes. The precipate was filtered and air dried to afford the product (10.5g, 93%).
¹H NMR (CDCl₃) 3.7 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 7

35 Preparation of methyl-5-chloro-2-(4-chlorophenyl)thiazol-4-ylacetic ester (Compound Number 14 in Table I).

5-Chloro-2-(4-chlorophenyl)thiazol-4-yl acetic acid (3.5g, 12mmol) prepared as described in Example 6, was dissolved in a solution of hot MeOH (50ml) and concentrated. H₂SO₄ (0.1ml) and the resulting reaction mixture heated under reflux for 4 hours. The reaction mixture was cooled to room temperature and poured into H₂O (50ml). The solution was neutralized (NaHCO₃) and the precipitate formed filtered and air dried to afford the product (3.0g, 83%).
¹H NMR (CDCl₃) 3.7 (3H, s, OCH₃), 3.8 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 8

50 Preparation of Ethyl-5-chloro-2-(4-chlorophenyl) thiazol-4-yl acetic ester (Compound No. 20 in Table I).

The reaction as described in Example 7 was carried out using ethanol instead of methanol to give the desired product (3.0g, 79%).
¹H NMR (CDCl₃) 1.15 (3H, t, CH₂CH₃), 3.8 (2H s, CH₂CO), 4.1 (2H, q, CH₂CH₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 9

Preparation of n-Propyl-5-chloro-2-(4-chlorophenyl) thiazol-4-yl acetic ester (Compound No. 15 in Table I).

The reaction as described in Example 7 was carried out using n-propanol instead of methanol to give the desired product (3.0g, 78%)


¹H NMR (CDCl₃) 0.95 (3H, t, CH₂CH₂CH₃), 1.7 (2H, hexet, CH₂CH₂CH₃), 3.85 (2H, s, CH₂CO), 4.1 (2H, t, CH₂CH₂CH₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 10

Preparation of Ethyl-2-(4-bromophenyl)-5,6-dihydrocyclopentadienyl thiazol-4-yl-carboxylic ester (Compound No. 192 in Table II).

A mixture of 5-bromo-2-carbethoxycyclopentane [described by A.M. Khaletskii et al, Zhur Obsheci, Khm. 31, 737 (1961); 7g 29.9mmol] and 4-bromothiobenzamide (3g, 14 mmol) were heated under reflux in ethanol (70ml) for 6 hours. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The residue was dissolved in ethyl acetate washed consecutively with aqueous NaOH (10%), brine, H₂O, dried (MgSO₄) filtered and concentrated under reduced pressure. The product was purified by flash chromatography (eluting with 10% ethyl acetate/Hexane) to afford the desired product (1.6g, 33%).

¹H NMR (CDCl₃) 1.30 (3H, t, OCH₂CH₃), 2.82 (2H, m, CH₂CO),



4.04 (1H, t, CH₂CO₂CH₂CH₃), 4.22 (2H, q, OCH₂CH₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 11

Preparation of Ethyl-2-phenyl-5,6-dihydrocyclopentadienylthiazol-4-yl carboxylic ester (Compound No. 183 in Table II).

A mixture of 5-bromo-2-carbethoxycyclopentane (14.5g, 0.62mmol) and thiobenzamide (7g, 0.05mmol) were heated under reflux in toluene (75ml) for 90 minutes. The toluene solution was decanted from the oil residue and organic layer washed with sodium bicarbonate solution, water, dried (MgSO₄), filtered and concentrated under reduced pressure. The product (0.99g, 6.5%) was purified by flash chromatography (eluting with 15% EtOAc/hexane).

¹H NMR (CDCl₃) 1.28 (3H, t, OCH₂CH₃), 2.8 (2H, t, ring CH₂), 2.98 and 3.10 (1H, m, ring CH₂), 4.04 (1H, t, CH CO₂Et), 4.22 (2H, q, OCH₂CH₃) and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 12

Preparation of Ethyl-2-(4- t-butylphenyl)-5,6-dihydrocyclopentadienylthiazol-4-ylcarboxylic ester (Compound No 184 in Table II)

The reaction as described in Example 11 was carried out using 4-t-butylthiobenzamide (6.0g, 0.03mol) instead of thiobenzamide to give the desired product (1.72g, 17.5%).
¹H NMR (CDCl₃) 1.32 (12H, m, ¹Bu and OCH₂CH₃), 2.9 (4H, m, ring H), 4.04 (1H, t, CH-CO₂Et), 4.24 (2H, q, OCH₂CH₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 13

Preparation of Ethyl-2-(4-trifluoromethylbenzene)-5,6-dihydrocyclopentadienylthiazol-4-yl-carboxylic ester
(Compound No. 185 in Table II).

The reaction as described in Example 10 was carried out using 4-trifluorothiobenzamide (6.0g, 0.029mol) instead of 4-bromothiobenzamide to give the desired product (2.47g, 25%).
¹H NMR (CDCl₃) 1.30 (3H, t, OCH₂CH₃), 2.82 (2H, m, ring CH₂), 2.98 and 3.15 (2H, m, ring CH₂), 4.05 (1H, t, CH CO₂Et), 4.25 (2H, q, OCH₂CH₃), and 7.85 (4H, AB quartet, aromatic C-H).

EXAMPLE 14

Preparation of Ethyl-2-(3-chloro-4-fluorophenyl)-5,6-dihydrocyclopentadienylthiazol-4-ylcarboxylic ester.
(Compound No. 188 in Table II).

The reaction as described in Example 10 was carried out using 3-chloro-4-fluorothiobenzamide (5g, 26mmol) instead of 4-bromothiobenzamide to give the desired product (2.55g, 29%).
¹H NMR (CDCl₃) 1.31 (3H, t, OCH₂CH₃), 2.83 (2H, m, ring CH₂), 3.0 and 3.10 (2H, m, ring CH₂), 4.03 (1H, t, CH CO₂Et), 4.23 (2H, q, OCH₂CH₃), 7.17 (1H, t, aromatic C-H), 7.74 (1H, m, aromatic C-H), and 7.99 (1H, d, aromatic C-H).

EXAMPLE 15

Preparation of Ethyl-2-(4-fluorophenyl)-5,6-dihydrocyclopentadienyl thiazol-4-yl-carboxylic ester. (Compound
No. 197 in Table II).

The reaction as described in Example 10 was carried out using 4-fluorothiobenzamide instead of 4-bromothiobenzamide to give the desired product (2.19g, 29%).
¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 2.8 (2H, t, ring CH₂), 2.9 and 3.1 (2H, m, ring CH₂), 4.0 (1H, t, CH CO₂Et), 4.2 (2H, q, OCH₂CH₃), and 7.5 (4H, AB quartet, aromatic C-H).

EXAMPLE 16

Preparation of Compound No. 59 in Table I

Bromine (3ml, 58 mmol) was added dropwise (2min) to ethyl-2-isopropylacetoacetate (10.4ml 58mmol) in ether (20ml) at 0°C (ice bath) and the resulting mixture stirred for a further 20 minutes. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in EtOH (70ml), 4-chlorothiobenzamide (5.1g) was added and the resulting solution refluxed for 5 hours. The reaction mixture was cooled, and concentrated under reduced pressure. The residue was partitioned between 10% NaOH and EtOAc, and the organic layer dried (MgSO₄), filtered and concentrated in vacuo. The product (3.7g,

38%) was purified by flash chromatography (eluted with 10% EtOAC/hexane).

¹H NMR (CDCl₃) 0.9 [3H, d, CH(CH₃)₂], 1.05 (3H, d, CH(CH₃)₂), 1.26 (3H, t, OCH₂CH₃), 2.45 (1H, m, CH-(CH₃)₂), 3.70 (1H, d, CHCO₂Et), 4.18 (2H, m, OCH₂CH₃), 7.25 (1H, s, thiazole ring H), and 7.6 (4H, AB, aromatic C-H).

EXAMPLE 17

Preparation of Compound No. 60 in Table I.

Bromine (0.5ml, 9.7mmol) in acetic acid (4.5ml) was added dropwise to compound 59 (3.16g 9.7mmol) in acetic acid (30ml) and the resulting mixture was heated at 90°C for 14 hours. The reaction mixture was cooled, then poured in saturated sodium bicarbonate solution and neutralised. The oil was extracted onto chloroform (3x) and the combined organic extracts dried (MgSO₄), filtered and concentrated under reduced pressure. The product was purified by flash chromatography (eluted with 2% EtOAC/Hexane) (0.9g, 25%).

¹H NMR (CDCl₃) 0.85 (3H, d, CH(CH₃)₂), 1.10 (3H, d, CH(CH₃)₂), 1.25 (3H, t, OCH₂CH₃), 2.72 (1H, m, CH-(CH₃)₂), 3.68 (1H, d, CHCO₂Et), 4.16 (2H, q, OCH₂CH₃), and 7.55 (4H, AB quartet, aromatic C-H).

EXAMPLE 18

Preparation of Compound No. 61 in Table I.

NaH (0.8g, 50% oil dispersion, 1.05 equiv) was added to compound 20 (5g, 15.8 mmol) dissolved in dry dimethoxyethane (DME) (35ml) and the resulting mixture stirred for 10 minutes at room temperature. Iodomethane (1.2 equiv) was added dropwise, and the resulting mixture was stirred for a further 2 hours. Water was added and the solution was extracted with chloroform (3x). The combined organic extracts were dried (MgSO₄), filtered, and concentrated under reduced pressure. The product was purified by chromatography (6% EtOAC/Hexane) (2.63g, 50%).

¹NMR (CDCl₃) 1.24 (3H, t, OCH₂CH₃), 1.60 (3H, d, CH(CH₃)CO₂Et), 4.05 (1H, q, CH(CH₃)CO₂Et) 4.17 (2H, q, OCH₂CH₃), and 7.5 (4H, AB quartet, aromatic C-H).

EXAMPLE 19

Preparation of N,N-Dimethyl-5-chloro-2-phenylthiazol-4-yl-acetamide (Compound No. 28 in Table I.)

A mixture of 5-chloro-2-phenylthiazol-4-yl-acetic acid (1.98g, 7.8mmol), tetramethylsulphurous diamide (1.01g, 9mmol) and chloroform (dry, 30ml) were heated under reflux for 4 hours. The reaction mixture was cooled and was concentrated under reduced pressure. The residue was taken up into chloroform (100ml) and H₂O (50ml) and an excess of sodium bicarbonate added. After filtration the chloroform layer was washed with dilute HCl, water, dried (MgSO₄), filtered and was concentrated under reduced pressure. The product was purified from hot petroleum ether (100-126°C) (0.47g) m.p. 77-79°C.

Microanalysis C: 55.62, H: 4.04; N: 9.98 (expected C: 55.55; H: 4.77; N: 9.97).

EXAMPLE 20

Preparation of N,N-Dimethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetamide (Compound No. 62 in Table I).

A mixture of 5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic acid (19.1g, 64mmol), thionyl chloride (1.2 equiv., 48ml 77mmol), and chloroform were heated at 80°C for 1 hour. The reaction mixture was cooled and then concentrated under reduced pressure. The residue was dissolved in chloroform (200ml) then saturated with dimethylamine (gas). The reaction mixture was heated under reflux for 2 hours, and concentrated under reduced pressure. The product was purified by flash chromatography (eluted with 70% EtOAc/Hexane) (14.0g, 66%).

¹H NMR (CDCl₃) 3.0 (3H, s, N(CH₃)₂), 3.2 (3H, s, N(CH₃)₂), 3.9 (2H, s, CH₂CON(CH₃)₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 21

Preparation of N,N-Dimethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetohydrazide. (Compound No. 63 in Table I).

The reaction as described in Example 20 was carried out using N,N-dimethylhydrazine (1.16ml) instead of dimethylamine to give the desired product.

¹H NMR (CDCl₃) 3.2 (6H, s, NHN(CH₃)₂), 3.8 (2H, s, CH₂CONHN(CH₃)₂), 7.6 (4H, AB quartet, aromatic C-H) and 12.8 (1H, bs, NHN(CH₃)₂).

EXAMPLE 22

Preparation of N-Butyl-5-bromo-2-(4-chlorophenyl)thiazol-4-ylacetic ester. (Compound No. 64 in Table 1).

The reaction as described in Example 7 was carried out using n-Butanol and refluxing for 9 hours instead of methanol and refluxing for 4 hours to give the desired product (1.1g, 47%).

¹H NMR (CDCl₃) 0.9 (3H, t, CH₃), 1.35 (2H, s, OCH₂CH₂CH₂CH₃), 1.6 (2H, s, OCH₂CH₂CH₂CH₃), 3.8 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 23

Preparation of Methylsulphonyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetamide. (Compound No. 68 in Table I).

The reaction as described in Example 20 was carried out using methylsulphonamide (1.45g, 15.2mmol) and 14 hours refluxing in place of dimethylamine and 2 hours refluxing. The precipitate was isolated by suction, washed with boiling MeOH, to give the desired product (0.7g).

¹H NMR (CDCl₃) 3.2 (3H, s, NHSO₂Me), 3.8 (2H, s, CH₂CO), 7.7 (4H, AB quartet, aromatic C-H) and 12.1 (1H, bs, NHSO₂Me).

EXAMPLE 24

Preparation of N-Phenyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetamide. (Compound No. 69 in Table I).

A mixture of 5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic acid (1g, 3 mmol), thionyl chloride (2 equiv., 0.72g, 6mmol), and chloroform were heated under reflux for 2.5 hours. The reaction mixture was cooled and concentrated under reduced pressure. The residue was dissolved in CHCl₃ (dry), followed by dropwise

addition of aniline (2 equiv., 0.56g, 6 mmol) and the resulting mixture refluxed for a further 5 hours. The reaction mixture cooled to room temperature and the precipitate was collected by filtration. The filtrate was concentrated under reduced pressure. The residue was triturated with EtOAc, and filtered to yield the desired product (0.4g, 32%).

¹H NMR (CDCl₃) 3.8 (2H, s, CH₂CONHC₆H₅), 7.0 (1H, dd, NHC₆H₅), 7.2 (2H, dd, NHC₆H₅), 7.6 (2H, d, NHC₆H₅), 7.4 (4H, AB quartet, aromatic C-H), and 10.2 (1H, s, HNC₆H₅).

EXAMPLE 25

Preparation of Ethyl-2-(4-iodophenyl)thiazol-4-yl acetic ester. (Compound No. 67 in Table I).

A mixture of 4-iodothiobenzamide (4.6g, 17.7mmol), toluene (dry 120ml), pTsOH (cat., 20mg), and ethyl-4-bromoacetoacetate were heated under reflux in a Dean and Stark apparatus for 4 hours. The reaction mixture was cooled and concentrated under reduced pressure, and the product purified by chromatography (eluted with 25% EtOAc/petrol; 2.6g, 39%).

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 3.9 (2H, s, CH₂COCH₂CH₃), 4.2 (2H, q, OCH₂CH₃), and 7.2 (1H, s, thiazolyl ring H), and 7.75 (4H, AB quartet, aromatic C-H).

EXAMPLE 26

Preparation of Ethyl-5-bromo-2-(4-iodophenyl)thiazol-4-ylacetic ester. (Compound No. 70 in Table I).

The reaction described in Example 1 was repeated using ethyl-2-(4-iodophenyl)thiazol-4-yl acetic ester (0.83g) in place of 2-(4-chlorophenyl)thiazol-4-yl acetic acid to yield the desired product (0.3g).

¹H NMR (CDCl₃) 1.15 (3H, t, OCH₂CH₃), 3.8 (2H, s, CH₂CO₂Et), 4.05 (2H, q, OCH₂CH₃), and 7.7 (4H, AB quartet, aromatic C-H).

EXAMPLE 27

Preparation 2-Hydroxyethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 72 in Table I).

The reaction described in Example 4 was repeated using ethyleneglycol (10ml, excess) in place of n-propanol to yield the desired product (0.52g, 46%).

¹H NMR (CDCl₃) 3.8 (2H, t, OCH₂CH₂OH), 3.9 (2H, s, CH₂CO), 4.3 (2H, t, OCH₂CH₂OH), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 28

Preparation of 2-Methoxyethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 71 in Table I).

The reaction described in Example 4 was repeated using 2-methoxyethanol (10ml) in place of n-propanol to yield the desired product (0.4g, 23%).

¹H NMR (CDCl₃) 3.4 (3H, s, OCH₂CH₂OMe), 3.6 (2H, t, OCH₂CH₂OMe), 3.9 (2H, s, CH₂CO), 4.3 (2H, t, OCH₂CH₂OMe), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 29

5 Preparation of 3-Methylbutyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 73 in Table I).

Dicyclohexylcarbodiimide (DCC) (1.1 equiv., 0.76g 3.8mmol) in CHCl_3 was added dropwise to 3-methylbutanol (2 equiv., 0.61g, 7.0mmol), DMAP (cat., 40mg), and 5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic acid (1g, 3.5mmol) suspended in CH_2Cl_2 (dry, 20ml) at 0° (ice bath) and the resulting mixture stirred
10 for 20 minutes. The reaction mixture was stirred for a further 2 hours at room temperature, then concentrated under reduced pressure. The product was purified by flash chromatography (eluted in 10% EtOAc/Hexane) (0.5g).

^1H NMR (CDCl_3) 0.9 (6H, d, $\text{CH}(\text{CH}_3)_2$), 1.55 (2H, q, $\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$), 1.7 (1H, m, $\text{CH}(\text{CH}_3)_2$), 4.2 (2H, t, COCH_2CH_2), and 7.6 (4H, A quartet, aromatic C-H).
15

EXAMPLE 30

20

Preparation of Allyl-5-chloro-2(4-chlorophenyl)thiazol -4-yl-acetic ester. (Compound No. 74 in Table I).

The reaction as described in Example 29 was carried out using allyl alcohol (2 equiv., 0.4g, 6.9mmol) in place of 3-methylbutanol to give the desired product (350mg, 27%).

^1H NMR (CDCl_3) 3.95 (2H, s, CH_2CO), 4.55 (2H, d, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.3 (2H, m, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.95 (1H, m, $\text{OCH}_2\text{CH}=\text{CH}_2$), and 7.6 (4H, AB quartet, aromatic C-H).

30

EXAMPLE 31

Preparation of 3-Pentyl-5-chloro-2(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 75 in Table I).

35

The reaction as described in Example 29 was carried out using 3-pentanol (2 equiv., 0.612g, 6.9mmol) in place of 3-methylbutanol to give the desired product (250mg, 25%).

^1H NMR (CDCl_3) 0.75 (6H, t, $\text{CH}(\text{CH}_2\text{CH}_3)_2$), 1.45 (4H, m, $\text{CH}(\text{CH}_2\text{CH}_3)_2$), 3.8 (2H, s, CH_2CO), 4.65 (1H, p, $\text{CH}(\text{CH}_2\text{CH}_3)_2$), and 7.7 (4H, AB quartet, aromatic C-H).
40

EXAMPLE 32

45

Preparation of Propargyl-5-chloro-2(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 76 in Table I).

The reaction as described in Example 29 was carried out using propargyl alcohol (2equiv., 0.38, 6.9mmol) in place of 3-methylbutanol to give the desired product (360mg).

^1H NMR (CDCl_3) 1.5 (1H, s, $\text{CH}_2\text{C}\text{CH}$), 3.9 (2H, s, CH_2CO), 4.75 (2H, s, $\text{CH}_2\text{C}\text{CH}$), and 7.6 (4H, AB quartet, aromatic C-H).
50

EXAMPLE 33

55

Preparation of n-Pentyl-5-chloro-2(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 77 in Table I).

The reaction described in Example 29 was carried out using n-pentanol (2equiv., 0.612g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.5g, 41%).

¹H NMR (CDCl₃) 0.9 (3H, d, CH₂CH₃), 1.5-1.7 (6H, m, aliphatic H), 3.05 (3H, d, CH₂CO), 4.2 (2H, t, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 34

Preparation of 3,7-Dimethyloctyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 78 in Table I).

The reaction as described in Example 29 was carried out using 3,7 dimethyloctanol to give the desired product (540mg).

¹H NMR (CDCl₃) 0.85-1.7 (19H, m, aliphatic H), 3.8 (2H, s, CH₂CO), 4.2 (2H, t, OCH₂) and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 35

Preparation of 2-Benzyloxyethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 79 in Table I).

The reaction as described in Example 29 was carried out using 2-benzyloxyethanol (2 equiv., 1.1g, 6.9mmol) in place of 3-methylbutanol to give the desired product (150mg).

¹H NMR (CDCl₃) 3.65 (2H, t, OCH₂CH₂O), 3.9 (2H, s, CH₂CO), 4.35 (2H, t, OCH₂CH₂O), 4.5 (2H, s, OCH₂CH₂OCH₂C₆H₅), 7.35 (m, aromatic C-H), and 7.5 (4H, AB quartet, aromatic C-H).

EXAMPLE 36

Preparation of Neopentyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 80 in Table I).

The reaction as described in Example 29 was carried out using neopentyl alcohol (2 equiv., 0.636g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.32g).

¹H NMR (CDCl₃) 0.9 (9H, s, (CH₃)₃), 3.84 (2H, s, CH₂CO), 3.88 (2H, s, OCH₂C(CH₃)₃), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 37

Preparation of 2-Methylpropyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 81 in Table I).

The reaction as described in Example 29 was carried out using 2-methylpropanol (0.53g, 6.9mmol), in place of 3-methylbutanol to give the desired product (0.46g).

¹H NMR (CDCl₃) 0.9 (6H, d, CH(CH₃)₂), 1.95 (1H, septet, CH(CH₃)₂), 3.85 (2H, s, CH₂CO) 3.95 (2H, d, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 38

Preparation of 2-Methylbutyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 82 in Table I).

The reaction as described in Example 29 was carried out using 2-methylbutanol (0.63g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.5g).

¹H NMR (CDCl₃) 0.9 (6H, d, CHCH₃), 1.2 (1H, m, CHCH₃), 1.4-1.7 (3H, m, aliphatic H), 3.8 (2H, s, CH₂CO), 4.0 (2H, m, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 39

Preparation of 1-Methylbutyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 83 in Table I).

The reaction as described in Example 29 was carried out using 1-methylbutanol (0.636g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.4g).

¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.25 (3H, d, CHCH₃), 1.5 (4H, m, CH₂CH₂CH₃), 3.8 (2H, s, CH₂CO), 5.0 (1H, sextet, O-CH(CH₃)CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 40

Preparation of 2,3-Dimethylpropyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 84 in Table I).

The reaction as described in Example 29 was carried out using 2,3-dimethylpropanol (0.61g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.36g).

¹H NMR (CDCl₃) 0.9 (6H, d, CH(CH₃)₂), 1.2 (3H, d, CHCH₃), 1.8 (1H, sextet, CH(CH₃)₂), 3.8 (2H, s, CH₂CO), 4.9 (1H, sextet, OCH(CH₃)₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 41

Preparation of 1-Methylpropyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 85 in Table I).

The reaction as described in Example 7 was carried out using 1-methylpropanol (0.517g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.34g).

¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.25 (3H, d, CH(CH₃)), 1.6 (2H, sextet, CH(CH₃)CH₂CH₃), 3.8 (2H, s, CH₂CO), 4.9 (1H, sextet, CH(CH₃)), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 42

Preparation of Cyclohexylmethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 86 in Table I).

EP 0 368 592 A1

The reaction as described in Example 29 was carried out using cyclohexanemethanol (0.79g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.45g).

¹H NMR (CDCl₃) 0.9-1.7 (11H, m, cyclohexane C-H), 3.8 (2H, s, CH₂CO), 4.0 (2H, d, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 43

Preparation of 2-Methoxyisopropyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 87 in Table I).

The reaction as described in Example 29 was carried out using 2-methoxyisopropanol (0.63g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.2g).

¹H NMR (CDCl₃) 1.3 (3H, d, CHCH₃), 3.35 (3H, s, OMe), 3.45 (2H, m, CH₂OMe), 3.85 (2H, s, CH₂CO), 5.15 (1H, sextet, OCH(CH₃)), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 44

Preparation of n-Hexyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 88 in Table I).

The reaction as described in Example 29 was carried using n-hexanol (0.71g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.45g).

¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.3-1.6 (8H, m, aliphatic H), 3.8 (2H, s, CH₂CO), 4.1 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 45

Preparation of 3-Methoxybutyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 89 in Table I).

The reaction as described in Example 29 was carried out using 3-methoxybutanol (0.73g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.3g).

¹H NMR (CDCl₃) 1.1 (3H, d, CHCH₃), 1.8 (2H, m, CH₂CH(OMe)CH₃), 3.25 (3H, s, OCH₃), 3.35 (1H, sextet, CH(CH₃)), 3.8 (2H, s, CH₂CO), 4.25 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 46

Preparation of Benzyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound 90 in Table I).

The reaction as described in Example 29 was carried out using benzyl alcohol (0.75g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.49g).

¹H NMR (CDCl₃) 3.9 (2H, s, CH₂CO), 5.2 (2H, s, OCH₂C₆H₅), 7.35 (5H, m, OCH₂C₆H₅), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 47

Preparation of p-Methoxyphenyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 91 in Table I).

The reaction was described in Example 29 was carried out using p-methoxyphenol (0.86g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.16g).
¹H NMR (CDCl₃) 3.8 (3H, s, OCH₃), 4.05 (2H, s, CH₂CO), 6.95 (4H, AB quartet, O-C₆H₄OMe), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 48

Preparation of n-Octyl-5-chloro-2-(4-chlorophenyl)thiayol-4-yl acetic ester. (Compound No. 92 in Table I).

The reaction as described in Example 29 was carried out using n-octanol (0.9g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.57g).
¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.3-1.6 (12H, m, aliphatic C-H), 3.8 (2H, s, CH₂CO), 4.15 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 49

Preparation of p-Nitrophenyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 93 in Table I).

The reaction as described in Example 29 was carried out using p-nitrophenol (0.96g, 6.9mmol) and refluxing for 4 hours in place of 3-methylbutanol and stirring at room temperature for 2 hours, to give the desired product (0.08g).
¹H NMR (CDCl₃) 4.1 (2H, s, CH₂CO), 7.6 (4H, AB quartet, aromatic C-H), and 7.8 (4H, AB quartet, O-C₆H₄NO₂).

EXAMPLE 50

Preparation of n-Decyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 94 in Table I).

The reaction as described in Example 29 was carried out using n-decanol (2 equiv), 1.1g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.59g).
¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.3-1.6 (16H, m, aliphatic C-H), 3.8 (2H, s, CH₂CO), 4.15 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 51

Preparation of Phenyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 95 in Table I).

The reaction described in Example 29 was carried out using phenol (2 equiv, 0.51g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.07g).
¹H NMR (CDCl₃) 4.1 (2H, s, CH₂CO), 7.3 (5H, m, phenyl C-H), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 52

5 Preparation of Neopentyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 96 in Table I.)

The reaction as described in Example 29 was carried out using 5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid (1g, 3.0mmol) and neopentanol (2 equiv, 0.53g, 6mmol) in place of 5-chloro-2-(4-chlorophenyl)-
 10 thiazol-4-yl acetic acid and 3-methylbutanol to give the desired product (1.2g).
¹H NMR (CDCl₃) 0.9 (9H, s, CH₂C(CH₃)₃), 3.85 (2H, s, CH₂CO), 3.9 (2H, s, CH₂C(CH₃)₃), and 7.6 (4H, AB quartet, aromatic C-H).

15 EXAMPLE 53

20 Preparation of 2-Methoxyisopropyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 97 in Table I.)

The reaction as described in Example 52 was carried out using 2-methoxyisopropanol (2 equiv., 0.54, 8.0mmol) in place of neopentanol to give the desired product (0.15g).
¹H NMR (CDCl₃) 1.2 (3H, d, CHCH₃), 3.3 (3H, s, OCH₃), 3.4 (2H, t, CH₂OCH₃), 3.8 (2H, s, CH₂CO), 5.1 (1H, m, OCHCH₃), and 7.5 (4H, AB quartet, aromatic C-H).
 25

EXAMPLE 54

30

Preparation of Phenyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 98 in Table I.)

The reaction as described in Example 52 was carried out using phenol (2 equiv, 0.56g, 6.0mmol) in place of neopentanol to give the desired product (0.13g).
¹H NMR (CDCl₃) 4.1 (2H, s, CH₂CO), 7.25 (5H, m, phenyl C-H), and 7.6 (4H, AB quartet, aromatic C-H).
 35

EXAMPLE 55

40

Preparation of Benzyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 99 in Table I.)

The reaction as described in Example 52 was carried out using benzylalcohol (2 equiv., 0.65g, 6.0mmol) in place of neopentanol to give the desired product (0.25g).
¹H NMR (CDCl₃) 4.05 (2H, s, CH₂CO), 5.25 (2H, s, OCH₂C₆H₅), 7.5 (5H, m, OCH₂C₆H₅), and 7.5 (4H, AB quartet, aromatic C-H).
 45

50

EXAMPLE 56

55 Preparation of Cyclohexylmethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 100 in Table I.)

The reaction as described in Example 52 was carried out using cyclohexylmethanol (2 equiv., 0.68g,

8.0mmol) in place of neopentanol to give the desired product (1.2g).

¹H NMR (CDCl₃) 1.3 (11H, m, aliphatic C-H), 3.85 (2H, s, CH₂CO), 3.96 (2H, d, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

5

EXAMPLE 57

10 Preparation of p-Methoxyphenyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 101 in Table I).

The reaction as described in Example 52 was carried out using p-methoxyphenyl (2 equiv., 0.75g, 6.0mmol) in place of neopentanol to give the desired product (0.08g).

15 ¹H NMR (CDCl₃) 3.8 (3H, s, OCH₃), 4.1 (2H, s, CH₂CO), 6.95 (4H, AB quartet, O-C₆H₄-OMe), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 58

20

Preparation of n-Octyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 102 in Table I).

25 The reaction as described in Example 52 was carried out using n-octanol (2 equiv., 0.78g, 6.0mmol) in place of neopentanol to give the desired product (1.2g).

¹H NMR (CDCl₃) 0.9 (3H, t, CH₂CH₃), 1.3-1.9 (12H, m, aliphatic C-H), 3.85 (2H, s, CH₂CO), 4.15 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

30

EXAMPLE 59

35 Preparation of t-Butyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl-acetic ester. (Compound No. 103 in Table I).

The reaction as described in Example 29 was carried out using 4-butanol (2 equiv., 0.5g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.08g).

¹H NMR (CDCl₃) 1.6 (9H, s, C(CH₃)₃), 3.9 (2H, s, CH₂CO), and 7.75 (4H, AB quartet aromatic C-H).

40

EXAMPLE 60

45

Preparation of (3-Chloro-2,2-dimethyl-1-propyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 104 in Table I).

The reaction as described in Example 29 was carried out using 3-chloro-2,2-dimethyl-1-propanol (1.1 equiv., 0.46g, 3.7mmol) in place of 3-methylbutanol to give the desired product.

50 ¹H NMR (CDCl₃) 0.95 (6H, s, (CH₃)₂), 3.35 (2H, s, CH₂Cl), 3.95 (2H, s, CH₂CO), 4.0 (2H, s, OCH₂), and 7.6 (4H, AB quartet, aromatic C-H).

55

EXAMPLE 61

Preparation of (Trimethylsilyl)methyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 105 in Table I).

The reaction as described in Example 29 was carried out using trimethylsilylmethanol (1.1 equiv) in place of 3-methylbutanol to give the desired product.

¹H NMR (CDCl₃) 0.0 (9H, s, Si(CH₃)₃), 3.8 (4H, s, CH₂CO and OCH₂Si(CH₃)₃), and 7.55 (4H, AB quartet aromatic C-H).

EXAMPLE 62

Preparation of (2-Chloromethyl)isopropyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 106 in Table I).

The reaction as described in Example 29 was carried out using 2-chloromethylisopropanol (2 equiv., 0.754g, 6.9mmol) and refluxing for 7 hours in place of 3-methylbutanol and stirring at room temperature for 2 hours to give the desired product (0.22g).

¹H NMR (CDCl₃) 1.5 (6H, s, (CH₃)₂), 3.75 (2H, s, CH₂Cl), 3.8 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 63

Preparation of [2-(Methoxyethoxy)]ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 107 in Table I).

The reaction as described in Example 29 was carried out using 2-(2-methoxyethoxy)ethanol (2 equiv., 0.83g 6.9mmol) and refluxing for 4 hours in place of 3-methylbutanol and stirring for 2 hours at room temperature to give the desired product (0.33g).

¹H NMR (CDCl₃) 3.3 (3H, s, OCH₃), 3.5 (2H, m, OCH₂CH₂OCH₃), 3.6 (2H, m, OCH₂CH₂OCH₃), 3.7 (2H, t, OCH₂CH₂), 3.9 (2H, s, CH₂CO), 4.3 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet aromatic C-H).

EXAMPLE 64

Preparation of Ethylglycolyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester (Compound No. 108 in Table I).

The reaction as described in Example 29 was carried out using ethylglycolate (2 equiv., 0.723g, 6.9mmol) and refluxing for 10 hours in place of 3-methylbutanol and stirring for 2 hours at room temperature to give the desired product (0.4g).

¹H NMR (CDCl₃) 1.2 (3H, t, OCH₂CH₃), 4.0 (2H, s, CH₂CO), 4.2 (2H, q, OCH₂CH₃), 4.7 (2H, s, OCH₂CO₂Et), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 65

Preparation of Tetrahydrofurfuryl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 109 in Table I).

The reaction as described in Example 29 was carried out using tetrahydrofurfuryl alcohol (2 equiv., 8.7g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.35g).
¹H NMR (CDCl₃) 1.95 (4H, m, aliphatic C-H), 3.9 (2H, m, -O-CH₂), 4.0 (2H, s, CH₂CO), 4.25 (3H, m, O-CH₂-CH-O), and 7.7 (4H, AB quartet, aromatic C-H).

5

EXAMPLE 66

10

Preparation of Ethylmandelyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 110 in Table I).

The reaction as described in Example 64 was carried out using ethylmandelate (2 equiv., 1.25g, 6.9mmol) in place of ethylglycolate to give the desired product.
¹H NMR (CDCl₃) 1.2 (3H, t, OCH₂CH₃), 4.0 (2H, s, CH₂CO), 4.25 (2H, m, OCH₂CH₃), 6.0 (1H, s, OCHC₆H₅), 7.35 (3H, m, phenyl-H), 7.50 (2H, m, phenyl-H), and 7.55 (4H, ABq, aromatic C-H).

20

EXAMPLE 67

Preparation of (α)-γ-Butyrolactonyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 111 in Table I).

The reaction as described in Example 52 was carried out using α-hydroxyγ-butyrolactone (2 equiv, 0.61g, 6.0mmol) and stirring at room temperature for 2 days in place of neopentanol and stirring at room temperature for 2 hours to give the desired product (0.13g).
¹H NMR (CDCl₃)

30

2.3 (1H, m, O-CH-C^H_H),

35

2.75 (1H, m, O-CH-C^H_H),

40

3.95 (2H, s, CH₂CO), 4.3 (1H, m, O-CH^H),

45

4.45 (1H, m, O-C^H_H),

5.5 (1H, t, O-CH) and 7.6 (4H, AB quartet, aromatic C-H).

50

EXAMPLE 68

55

Preparation of Methylactatyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 112 in Table I).

EP 0 368 592 A1

The reaction as described in Example 64 was carried out using 5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid and methyl lactate (2 equiv., 0.723g, 6.9 mmol in place of 5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic acid and ether glycolate to give the desired product (0.4g).

¹H NMR (CDCl₃) 1.5 (3H, d, CH(CH₃)), 3.75 (3H, s, OCH₃), 3.95 (2H, s, CH₂CO), 5.2 (1H, q, CH(CH₃)), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 69

Preparation of Ethylmandelyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 113 in Table I).

The reaction as described in Example 67 was carried out using ethyl mandelate (2 equiv., 1.08g, 6.0mmol) in the place of α -hydroxy- γ -butyrolactone to give the desired product (0.36g).

¹H NMR (CDCl₃) 1.2 (3H, t, OCH₂CH₃), 4.0 (2H, s, CH₂CO), 4.2 (2H, m, OCH₂CH₃), 6.0 (1H, s, O-CH), 7.35 (3H, m, phenyl-H), 7.50 (2H, m, phenyl-H), and 7.55 (4H, AB quartet, aromatic C-H).

EXAMPLE 70

Preparation of Ethylglycolyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 114 in Table I).

The reaction as described in Example 67 was carried out using ethylglycolate (2 equiv, 8.626g, 6.0mmol) in place of α -hydroxy- γ -butyrolactone to give the desired product (0.2g).

¹H NMR (CDCl₃) 1.25 (3H, t, OCH₂CH₃), 4.0 (2H, s, CH₂CO), 4.2 (2H, q, OCH₂CH₃), 4.7 (2H s, OCH₂CO₂Et), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 71

Preparation of (2-Methoxyethoxy)ethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid. (Compound No. 115 in Table I).

The reaction as described in Example 67 was carried out using 2-methoxyethoxyethanol (2 equiv., 0.72g, 6.0mmol) in place of α -hydroxy- γ -butyrolactone to give the desired product (0.46g).

¹H NMR 3.35 (3H, s, OCH₃), 3.55 (4H, m, OCH₂CH₂OCH₃), 3.7 (2H, t, OCH₂CH₂O), 3.9 (2H, s, CH₂CO), 4.3 (2H, t, OCH₂CH₂O), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 72

Preparation of Methylmandelyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acid ester. (Compound No. 116 in Table I).

The reaction as described in Example 67 was carried out using methyl mandelate (2 equiv, 0.725g, 6.0mmol) in place of α -hydroxy- γ -butyrolactone to give the desired product (0.46g).

¹H NMR (CDCl₃) 3.7 (3H, s, OCH₃), 4.0 (2H, s, CH₂CO), 6.0 (1H, s, OCH), 7.35 (3H, m, phenyl C-H), 7.5 (2H, m, phenyl C-H), and 7.55 (4H, AB quartet, aromatic C-H).

EXAMPLE 73

5 Preparation of Methylmandelyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 117 in Table I).

The reaction as described in Example 64 was carried out using methylmandelate (2 equiv, 1.15g 6.9mmol) in place of ethyl glycolate to give the desired product (0.32g).

10 ¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 4.0 (2H, s, CH₂CO), 6.0 (1H, s, OCH), 7.35 (3H, m, phenyl C-H), 7.5 (2H, m, phenyl C-H and 7.55 (4H, AB quartet, aromatic C-H).

EXAMPLE 74

15

Preparation of (3,3,2,2-Tetrafluoro-2-methyl-2-butyl)-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 118 in Table I).

20

The reaction as described in Example 64 was carried out using 3,3,2,2-tetrafluoro-2-methyl-2-butanol (2 equiv, 1.11g, 6.9mmol) in place of ethylglycolate to give the desired product (0.5g).

¹H NMR (CDCl₃) 1.28 (3H, s, CH₃), 1.32 (3H, s, CH₃), 3.8 (2H, s, CH₂CO), 5.9 (1H, m, CHF₂), and 7.6 (4H, AB quartet, aromatic C-H).

25

EXAMPLE 75

30

Preparation of Methyl S(-)-lactatyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 119 in Table I).

The reaction as described in Example 64 was carried out using methyl-S(-)-lactate (2 equiv., 1.15g, 6.9mmol) in place of ethylglycolate to give the desired product (0.4g).

35 ¹H NMR (CDCl₃) 1.5 (3H, d, CH(CH₃)), 3.7 (3H, s, OCH₃), 3.9 (2H, s, CH₂CO), 5.2 (1H, q, CH(CH₃)), and 7.6 (4H, AB quartet, aromatic C-H).

40

EXAMPLE 76

Preparation of Ethylglycyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 120 in Table I).

45

The reaction as described in Example 29 was carried out using ethylglycine (2 equiv., 0.9g) and stirring at room temperature for 2 days in place of 3-methyl butanol and stirring at room temperature for 2 hours to give the desired product (0.06g).

50 ¹H NMR (CDCl₃) 1.2 (3H, t, OCH₂CH₃), 3.8 (2H, s, CH₂CO), 4.1 (2H, s, CH₂CO), 4.2 (2H, quartet, OCH₂CH₃), 7.5 (1H, bs, HNCH₂CO₂Et), and 7.7 (4H, AB quartet, aromatic C-H).

EXAMPLE 77


55

Preparation of ()-butyrolactonyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 121 in Table I).

Table I).

The reaction as described in Example 29 was carried out using α -hydroxy- γ -butyrolactone (2 equiv., 0.7g, 6.9mmol) in place of 3-methylbutanol to give the desired product (0.38g).

¹H NMR (CDCl₃)



3.95 (2H, s, CH₂CO), 4.4 (2H, m, O-CH₂), 5.5 (1H, t, OCH), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 78

Preparation of Ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl bromoacetic ester. (Compound No. 122 in Table I).

Lithium hexamethyldisilazane (LHMDS) (1 equiv., 1M solution in THF, 6.32ml, 6.3mmol) was added dropwise to ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester dissolved in dry THF (50ml) at -70°C under N₂. The reaction mixture was allowed to stir for 10 minutes followed by the addition of NBS (1 equiv., 1.12g, 6.3mmol) in THF (10ml) and then allowed to warm to room temperature. The precipitate was removed by suction and the filtrate was concentrated under reduced pressure and the product was purified by flash chromatography (eluted with 10% Et₂O/Hexane) (1.7g).

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 4.2 (2H, q, OCH₂CH₃), 5.7 (1H, s, CHBr) and 7.6 (4H, AB quartet, aromatic C-H)

EXAMPLE 79

Preparation of Compound No. 123 in Table I.

LHMDS (1 equiv., 0.26g, 1.57mmol) was added dropwise to ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl acetic ester dissolved in THF (7ml) at -70°C (under N₂) and the resulting mixture stirred for 15 minutes. Benzylchloroformate (1 equiv., 0.27g, 0.23, 1.57mmol) was added and the reaction mixture was allowed to warm to room temperature and stirred for 48 hours. Sat NH₄Cl solution was added and the organic layer separated, and concentrated under reduced pressure. The product (0.25g) was purified by chromatography (eluted with 10% Et₂O/Hexane).

¹H NMR (CDCl₃) 1.05 (3H, t, OCH₂CH₃), 4.05 (2H, quartet, OCH₂CH₃), 4.8 (1H, s, CHCO₂CH₂C₆H₅), 5.05 (2H, s, OCH₂C₆H₅), 7.1 (5H, m, phenyl C-H), and 7.3 (4H, AB quartet, aromatic C-H).

EXAMPLE 80

Preparation of Ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl chloroacetic ester. (Compound No. 124 in Table I).

The reaction as described in Example 78 was carried out using NCS (1 equiv., 0.21g, 1.57mmol), in place of NBS to give the desired product (0.3g).
¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 4.3 (2H, q, OCH₂CH₃), 5.7 (1H, s, CHCl), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 81

Preparation of Compound No. 125 in Table I).

A mixture of Cu(I)CN (1 equiv., 0.115g, 1.57mmol), N-methylpyrrolidinone (5ml) and the thiazole derivative (compound No. 61) (500mg, 1.57mmol) were heated at 140°C for 10 minutes. The reaction mixture was cooled and the solvent distilled off (approx 44° under high vac.) and the residue partitioned between CHCl₃ and H₂O. The insoluble salts were filtered off under suction, the organic layer separated, dried (MgSO₄), filtered and concentrated under reduced pressure. The product (0.1g) was purified by chromatography (eluted with 10% Et₂O/Hexane).

¹H NMR (CDCl₃) 1.45 (3H, t, OCH₂CH₃), 4.5 (2H, q, OCH₂CH₃) and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 82

Preparation of Methyl-5-nitro-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 126 in Table I).

Methyl-2-(4-chlorophenyl)thiazol-4-yl acetic ester (5.0g, 18.7mmol) in CH₃CN (dry, 10ml) was carefully added dropwise to BF₄NO₂ (0.5equiv., 1.25g, 9.3mmol) suspended in CH₃CN (50ml) at 5°C (ice/salt bath). When all the BF₄NO₂ salt had dissolved, H₂O (10ml) was added, and the mixture concentrated under reduced pressure to approximately 20ml volume. The solution was extracted with CHCl₃ and the combined organic extracts dried (MgSO₄), filtered and concentrated under reduced pressure. The product (1.5g) was purified by chromatography (eluted with 25% Et₂O/Hexane).

¹H NMR (CDCl₃) 3.8 (3H, s, OCH₃), 4.3 (2H, s, CH₂CO) and 7.7 (4H, AB quartet, aromatic C-H).

EXAMPLE 83

Preparation of Methyl-5-bromo-2-(4-fluorophenyl)thiazol-4-yl acetic ester. (Compound No. 127 in Table I).

Bromine (1.6g, 0.52ml, 10mmol) was added dropwise to 2-(4-fluorophenyl)thiazol-4-yl acetic acid (2.37g, 10mmol) dissolved in AcOH (30ml) and the resulting mixture stirred at room temperature for 2 hours. The precipitate was separated by suction, washed with CH₂Cl₂ and air dried to yield the corresponding 5-bromo thiazole acid derivative. This was esterified in the usual way to give the desired product (2.0g).

¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 3.9 (2H, s, CH₂CO), 7.1 (2H, t, aromatic C-H), and 7.85 (2H, dd, aromatic C-H).

EXAMPLE 84

Preparation of Ethyl-5-bromo-2-(4-fluorophenyl)thiazol-4-yl acetic ester. (Compound No. 128 in Table I).

5-bromo-2-(4-fluorophenyl)thiazol-4-yl acetic acid (0.5g, 1.58mmol) (preparation outlined in Example 83) was esterified in the usual way (see Example 3) to give the desired product (0.46g).

¹H NMR (CDCl₃) 1.28 (3H, t, OCH₂CH₃), 3.85 (2H, s, CH₂CO), 4.25 (2H, q, OCH₂CH₃), 7.12 (2H, m, aromatic C-H), and 7.8 (2H, m, aromatic C-H).

EXAMPLE 85

Preparation of n-Pentyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 129 in Table I).

The reaction as described in Example 52 was carried out using n-pentanol (2 equiv, 0.53g, 6mmol) in place of neopentanol to give the desired product (0.4g).

¹H NMR (CDCl₃) 0.85 (3H, t, CH₂CH₃), 1.3 (4H, m, aliphatic H's), 1.63 (2H, m, aliphatic H), 3.85 (2H, s, CH₂CO), 4.15 (2H, t, OCH₂CH₂), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 86

Preparation of n-Hexyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 130 in Table 1).

The reaction as described in Example 52 was carried out using n-hexanol (2 equiv, 0.614g, 6.0mmol), in place of neopentanol to give the desired product (0.35g).

¹H NMR (CDCl₃) 1.0 (3H, t, CH₂CH₃), 1.45 (8H, m, aliphatic H), 1.8 (2H, m, aliphatic H), 4.0 (2H, s, CH₂CO), 4.3 (2H, t, OCH₂CH₂), and 7.75 (4H, AB quartet, aromatic C-H)

EXAMPLE 87

Preparation of Ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl fluoroacetic ester. (Compound No. 132 in Table I).

A mixture of ethyl-5-chloro-2-(4-chlorophenyl)thiazol-4-yl bromoacetic ester (Compound No. 122) (1g, 2.5mmol), CsF (3 equiv., 1.15g, 7.5mmol) and dry DMF (10ml) were heated at 130°C for 1 hour. The reaction mixture was cooled and then concentrated under reduced pressure. The product (0.15g) was purified by chromatography (eluted with 20% Et₂O/Hexane).

¹H NMR (CDCl₃) 1.35 (3H, t, OCH₂CH₃), 4.35 (2H, q, OCH₂CH₃), 6.05 (1H, d, CHF) and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 88

Preparation of Methyl-5-nitro-2-(4-fluorophenyl)thiazol-4-yl acetic ester. (Compound No. 137 in Table I).

The reaction as described in Example 82 was carried out using methyl-2-(4-fluorophenyl)thiazol-4-yl acetic ester (Compound No. 139) (1.5g, 5.6mmol) in place of methyl-2-(4-chlorophenyl)thiazol-4-yl acetic ester to give the desired product (0.2g).

¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 4.3 (2H, s, CH₂CO), 7.2 (2H, dd, aromatic C-H), and 8.0 (2H, dd, aromatic C-H).

EXAMPLE 89

Preparation of 5-Bromo-2-(3-chlorophenyl)thiazol-4-yl acetic acid. (Compound No. 140 in Table I).

The reaction as described in Example 1 was carried out using 2-(3-chlorophenyl)thiazol-4-yl acetic acid in place of 2-(4-chlorophenyl)thiazol-4-yl acetic acid to give the desired product.

¹H NMR (d₆-DMSO) 3.92 (2H, s, CH₂CO), 7.4 (2H, m, aromatic C-H), 7.70 (1H, m, aromatic C-H), and 7.87 (1H, d, aromatic C-H).

EXAMPLE 90

Preparation of Ethyl-5-bromo-2-phenylthiazol-4-yl acetic ester. (Compound No. 141 in Table I).

5-bromo-2-phenylthiazol-4-yl acetic acid, (obtained by simple bromination on the corresponding acid, see for eg, Example 83) (0.5g, 1.68mmol) was esterified in the usual way to give the desired product (0.13g).

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 3.85 (2H, s, CH₂CO), 4.2 (2H, q, OCH₂CH₃), 7.4 (3H, m, aromatic C-H), and 7.85 (2H, m, aromatic C-H).

EXAMPLE 91

Preparation of Methyl-5-bromo-2-phenylthiazol-4-yl acetic ester. (Compound No. 142 in Table I).

The reaction as described in Example 3 was carried out using 5-bromo-2-phenylthiazol-4-yl acetic acid (0.5g, 1.68mmol) in place of 5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid to give the desired product (0.17g).

¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 3.9 (2H, s, CH₂CO), 7.4 (3H, m, aromatic C-H), and 7.85 (2H, m, aromatic C-H).

EXAMPLE 92

Preparation of Ethyl-5-bromo-2-(3-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 143 in Table I).

5-bromo-2-(3-chlorophenyl)thiazol-4-yl acetic acid (0.4g, 1.2mmol) was esterified in the usual way to give the desired product (0.25g).

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 3.85 (2H, s, CH₂CO), 4.2 (2H, q, OCH₂CH₃), 7.4 (2H, m, aromatic C-H), 7.7 (1H, m, aromatic C-H), and 7.9 (1H, m, aromatic C-H).

EXAMPLE 93

Preparation of Methyl-5-bromo-2-(3-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 144 in Table I).

The compound was prepared by methods analogous to that described in Example 92.

¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 3.9 (2H, s, CH₂CO), 7.4 (2H, m, aromatic C-H), 7.7 (1H, m, aromatic C-H), and 7.9 (1H, d, aromatic C-H).

EXAMPLE 94

Preparation of Methyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl bromoacetic ester. (Compound No. 145 in Table I).

A mixture of 2-(4-chlorophenyl)thiazol-4-ylacetic acid (8.0g, 31.6mmol), bromine (2ml) and acetic acid (100ml) were heated at 100°C for 1.5 hours. The reaction mixture was cooled, and the precipitate separated by suction. Half of the solid was refluxed in methanol for 5 hours. The reaction mixture was concentrated and the product (1.1g) purified by chromatography.
¹H NMR (CDCl₃) 3.85 (3H, s, OCH₃), 5.75 (1H, s, CHBr), and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 95

Preparation of Ethyl-5-bromo-(4-chlorophenyl)thiazol-4-yl bromoacetic ester. (Compound No. 146 in Table I).

The remaining half of the solid from Example 94 was refluxed in ethanol for 5 hours. The reaction mixture was cooled, filtered and the filtrate concentrated under reduced pressure. The product (1.0g) was purified by chromatography.
¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 4.2 (2H, q, OCH₂CH₃), 5.7 (1H, s, CHBr), and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 96

Preparation of methyl-5-bromo-2-(4-bromophenyl)thiazol-4-yl acetic ester. (Compound No. 147 in Table I).

This compound was prepared by methods analogous to those described in Example 2.
¹H NMR (CDCl₃) 3.75 (3H, s, OCH₃), 3.9 (2H, s, CH₂CO), and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 97

Preparation of Ethyl-5-bromo-2-(4-bromophenyl)thiazol-4-yl acetic ester. (Compound No. 148 in Table I).

This compound was prepared by methods analogous to those described in Example 3.
¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 3.85 (CH₂CO), 4.2 (2H, q, OCH₂CH₃), and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 98

Preparation of n-Propyl-5-bromo-2-(4-bromophenyl)thiazol-4-yl acetic ester. (Compound No. 149 in Table I).

This compound was prepared by methods analogous to those described in Example 7.
¹H NMR (CDCl₃) 0.95 (3H, s, OCH₂CH₂CH₃), 1.7 (2H, sextet, OCH₂CH₂CH₃), 3.85 (2H, s, CH₂CO), 4.1 (2H, t, OCH₂CH₂CH₃), and 7.65 (4H, AB quartet, aromatic C-H).

EXAMPLE 99

5 Preparation of 5-Bromo-2-(4-bromophenyl)thiazol-4-yl acetic acid. (Compound No. 150 in Table I).

The reaction as described in Example 83 was carried out using 2-(4-bromophenyl)thiazol-4-yl acetic acid (4g, 13.5mmol) to give the desired product (3.76g).

¹H NMR (D₆-DMSO + CDCl₃) 3.75 (2H, s, CH₂CO), 7.8 (4H, AB quartet, aromatic C-H).

10

EXAMPLE 100

15

Preparation of 5-Bromo-2-(4-trifluoromethylphenyl)thiazol-4-yl acetic acid. (Compound No. 151 in Table I).

This compound was prepared by methods analogous to those described in Example 83.

¹H NMR (D₆-DMSO + CDCl₃) 3.9 (2H, s, CH₂CO₂H), and 7.85 (4H, AB quartet, aromatic C-H).

20

EXAMPLE 101

25

Preparation of Methyl-5-bromo-2-(4-trifluoromethylphenyl)thiazol-4-yl acetic ester. (Compound No. 152 in Table I).

This compound was prepared by a reaction analogous to that described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.75 (3H, s, CO₂CH₃), 3.9 (2H, s, CH₂CO), and 7.8 (4H, AB quartet, aromatic C-H).

35

EXAMPLE 102

Preparation of Ethyl-5-bromo-2-(4-trifluoromethylphenyl)thiazol-4-yl acetic ester. (Compound No. 153 in Table I).

This compound was prepared by methods analogous to those described in Example 83 followed by esterification as described in Example 3.

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 3.9 (2H, s, CH₂CO), 4.2 (2H, q, OCH₂CH₃), and 7.8 (4H, AB quartet, aromatic C-H).

45

EXAMPLE 103

50

Preparation of Methyl-5-amino-2-(4-chlorophenyl)thiazol-4-yl acetic ester. (Compound No. 154 in Table I).

Methyl-5-nitro-2-(4-chlorophenyl)thiazol-4-yl acetic ester (Compound No. 126) (0.85g) in MeOH/CHCl₃ - (50ml, 5ml) was hydrogenated over 10% palladium on charcoal (cat). The product (0.2g) was purified by chromatography (eluted with 20% Et₂O/Hexane). ¹H NMR (CDCl₃) 3.7 (3H, s, CO₂CH₃), 3.8 (2H, s, CH₂CO), and 7.5 (4H, AB quartet, aromatic C-H).

55

EXAMPLE 104

5 Preparation of 5-Bromo-2-(2,4-difluorophenyl)thiazol-4-yl acetic acid. (Compound No. 155 in Table I).

This compound was prepared by methods analogous to those described in Example 83.

¹H NMR (CDCl₃) 3.75 (2H, s, CH₂CO), 7.53 (1H, m, aromatic C-H), 7.74 (7H, m, aromatic C-H), 7.9 (1H, m, aromatic C-H) and 12.5 (1H, bs, CO₂H).

10

EXAMPLE 105

15

Preparation of Methyl-5-bromo-2-(2,4-difluorophenyl)thiazol-4-yl acetic ester. (Compound No. 157 in Table I).

20 This compound was prepared by methods analogous to that described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.8 (3H, s, CO₂CH₃), 4.1 (2H, s, CH₂CO), 7.0 (1H, m, aromatic C-H), 7.1 (1H, m, aromatic C-H), and 8.65 (1H, m, aromatic C-H).

25

EXAMPLE 106

30 Preparation of Ethyl-5-bromo-2-(2,4-difluorophenyl)thiazol-4-yl acetic ester. (Compound No. 158 in Table I).

This compound was prepared by a method as described in Example 83 followed by esterification as described in Example 3.

¹H NMR (CDCl₃) 1.3 (3H, t, OCH₂CH₃), 4.15 (2H, s, CH₂CO), 4.25 (2H, q, OCH₂CH₃), 7.0 (1H, m, aromatic C-H), 7.12 (1H, m, aromatic C-H), and 8.7 (1H, m, aromatic C-H).

35

EXAMPLE 107

40

Preparation of Methyl-5-bromo-2-(3,4-difluorophenyl)thiazol-4-yl acetic ester. (Compound No. 159 in Table I).

45 This compound was prepared as described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.75 (3H, s, CO₂CH₃), 3.88 (2H, s, CH₂CO), 7.25 (1H, m, aromatic C-H), 7.6 (1H, m, aromatic C-H), and 7.75 (1H, m, aromatic C-H).

50

EXAMPLE 108

55 Preparation of Ethyl-5-bromo-2-(3,4-difluorophenyl)thiazol-4-yl acetic ester. (Compound No. 160 in Table I).

This compound was prepared as described in Example 83 followed by esterification as described in Example 3.

¹H NMR (CDCl₃ + d₆DMSO)

1.33 (3H, t, OCH₂CH₃), 4.28 (2H, q, OCH₂CH₃), 4.32 (2H, s, CH₂CO), 7.4 (1H, m, aromatic C-H), and 8.15 (2H, m, aromatic C-H).

EXAMPLE 109

Preparation of 5-Bromo-2-phenylthiazol-4-yl acetic acid. (Compound No. 162 in Table I).

This compound was prepared by a method as described in Example 83.

¹H NMR (CDCl₃ + d₆DMSO)

3.85 (2H, s, CH₂CO), 7.45 (3H, m, aromatic C-H), and 7.89 (2H, m, aromatic C-H).

EXAMPLE 110

Preparation of 5-Bromo-2-(4-fluorophenyl)thiazol-4-yl acetic acid. (Compound No. 163 in Table I).

This compound was prepared by methods analogous to that described in Example 83.

¹H NMR (CDCl₃ + d₆DMSO)

3.84 (2H, s, CH₂CO), 7.13 (1H, t, aromatic C-H), 7.8 (1H, m, aromatic C-H), and 8.05 (1H, m, aromatic C-H).

EXAMPLE 111

Preparation of N,N-Dimethyl-5-bromo-2-(2,6-difluorophenyl)thiazol-4-yl acetamide. (Compound No. 164 in Table I).

The reaction as described in Example 83 was carried out using N,N-dimethyl-2-(2,6-difluorophenyl)thiazol-4-yl acetamide (1g, 3.5mmol) to give the desired product (1.1g).

¹H NMR (d₆DMSO) 2.95 (3H, s, NCH₃), 3.2 (3H, s, NCH₃), 4.0 (2H, s, CH₂CO), 7.4 (2H, m, aromatic C-H), and 7.7 (1H, m, aromatic C-H).

EXAMPLE 112

Preparation of N,N-Dimethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetohydrazide. (Compound No. 165 in Table I).

The reaction as described in Example 52 was carried out using N,N-dimethylhydrazine (0.28g, 4.84mmol) in place of neopentanol to give the desired product (recrystallised from EtOAC) (1.08g).

¹H NMR (CDCl₃) 2.55 (3H, s, NCH₃), 2.60 (3H, s, NCH₃), 4.0 (2H, s, CH₂CO), 6.1 (1H, bs, HNN (CH₃)₂), and 7.65 (4H, AB quartet aromatic (C-H)).

EXAMPLE 113

Preparation of Compound No. 166 in Table 1

Iodomethane (2ml excess) was added dropwise to N,N-dimethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetohydrazide (Example 112), (0.52g, 1.4mmol) suspended in dry methanol (15ml) and the result solution stirred at room temperature for 24 hours. The reaction mixture was concentrated under reduced pressure, and the residue triturated with Et₂O to give the desired product (0.14g).

¹H NMR (d₆ DMSO) 3.9 (9H, s, N(CH₃)₃), 3.95 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 114

Preparation of Methyl-5-bromo-2-(4-toluy)lthiazol-4-yl acetic ester. (Compound No. 167 in Table I).

This compound was prepared by a method as described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.4 (3H, s, aromatic CH₃), 3.75 (3H, s, CO₂CH₃), 3.90 (2H, s, CH₂CO), and 7.5 (4H, AB quartet, aromatic C-H).

EXAMPLE 115

Preparation of Ethyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl thioacetic ester. (Compound No. 168 in Table I).

A mixture of 5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic acid (0.5g, 1.5mmol), EtSH (2 equiv., 0.245g, 3.0mmol), pyridine (3 equiv., 0.625g) and phenyl dichlorophosphate (1.5equiv, 0.63g) was stirred at room temperature for 8 hours. The mixture was poured into iced water and extracted with CHCl₃. The combined organic extracts were dried (MgSO₄), filtered and concentrated under reduced pressure. The product (0.17g) was purified by chromatography (eluted with 5% Et₂O/Hexane).

¹H NMR (CDCl₃) 1.3 (3H, t, SCH₂CH₃), 2.9 (2H, q, SCH₂CH₃), 4.05 (2H, s, CH₂CO), and 7.65 (4H, AB quartet aromatic C-H).

EXAMPLE 116

Preparation of Methyl-5-bromo-2-(4-methoxyphenyl)thiazol-4-yl acetic ester. (Compound No. 169 in Table I).

This compound was prepared by a method as described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.75 (2H, s, CH₂CO), 3.85 (6H, s, CO₂CH₃), 7.4 (4H, AB quartet, aromatic C-H).

EXAMPLE 117

Preparation of 5-Bromo-2-(4-methoxyphenyl)thiazol-4-yl acetic acid. (Compound No. 170 in Table I).

This compound was prepared by a method as described in Example 83

¹H NMR (d₆ DMSO) 3.85 (2H, s, CH₂CO₂H), 3.87 (3H, s, OCH₃), and 7.45 (4H, AB quartet, C-H).

EXAMPLE 118

Preparation of Methyl-5-bromo-2-(4-trifluoromethoxy phenyl) thiazol-4-yl acetic ester. (Compound No. 171 in Table I).

This compound was prepared by a method as described in Example 83 followed by esterification as described in Example 2.

¹H NMR (CDCl₃) 3.75 (3H, s, CO₂CH₃), 3.9 (2H, s, CH₂CO), and 7.6 (4H, AB quartet, aromatic C-H).

EXAMPLE 119

Preparation of 5-Bromo-2-(4-acetylphenyl)thiazol-4-yl acetic acid. (Compound No. 172 in Table I).

This compound was prepared by a method as described in Example 83.

¹H NMR (d₆-DMSO) 2.6 (3H, s, COCH₃), 3.83 (2H, s, CH₂CO), and 8.0 (4H, m, aromatic C-H).

EXAMPLE 120

Preparation of Methyl-2-(4-chlorophenyl)thiazol-4-yl fluoroacetic ester. (Compound No. 173 in Table I).

A mixture of methyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester (0.5g, 1.4mmol), Ag(I)F (2 equiv., 0.35g, 2.8mmol), and CH₃CN (dry, 10ml) were stirred at room temperature for 24 hours. The reaction mixture was filtered and filtrate concentrated under reduced pressure to give the desired product (0.22g).

¹H NMR (CDCl₃) 3.9 (3H, s, CO₂CH₃), 6.0 (1H, d, CHF), 5.5 (1H, s, thiazole ring H), and 7.7 (4H, AB quartet aromatic C-H).

EXAMPLE 121

35

Preparation of Methyl-5-bromo-2-(4-hydroxyphenyl)thiazol-4-yl acetic ester. (Compound No. 174 in Table I).

This compound was prepared by a method as described in Example 83 followed by esterification in Example 2.

¹H NMR (CDCl₃ + d₆ DMSO)

3.7 (3H, s, CO₂CH₃), 3.8 (2H, s, CH₂CO), 7.3 (4H, AB quartet, aromatic C-H).

EXAMPLE 122

45

Preparation of Compound No. 176 in Table I.

50

A mixture of 4-chlorothiobenzamide (8.6g, 54.7mmol) dimethyl-3-oxo-2-chloroglutarate (Bader, 1 equiv, 10.2g) and dry methanol (100ml) were heated under reflux for 8 hours. The reaction mixture was cooled to room temperature and the the product (compound number 175) (8.6g) was obtained on filtration.

¹H NMR (CDCl₃) 3.70 (3H, s, CO₂CH₃), 3.9 (3H, s, CO₂CH₃), 4.3 (2H, s, CHCO), and 7.7 (4H, AB quartet, aromatic C-H).

Compound No. 175 (Example 122) (2.0g) was base hydrolysed in the usual way to give the desired product (1.3g).

¹H NMR (d₆ DMSO) 4.25 (2H, s, CH₂CO), and 7.7 (4H, AB quartet, aromatic C-H).

EXAMPLE 124

5 Preparation of 5-Hydroxy-2-(4-chlorophenyl)thiazol-4-ylacetic acid. (Compound No. 177 in Table I).

A mixture of methyl-4-chlorothiobenzoate (2.11g, 11.3mmol), aspartic acid (2.7g, 2.02mmol), aq NaOH (3N, 8.8ml), and Et₂O (8ml) were vigorously stirred at room temperature for 3 days. The aq. layer was separated, acidified (dil.HCl), and extracted with Et₂O (2x25ml). The combined ether extracts were dried
 10 (MgSO₄), filtered, and concentrated under reduced pressure to yield N-p-chlorothiobenzoylaspartic acid (0.23g).

¹H NMR (CDCl₃ + d₆ DMSO)

3.4 (2H, dq, CH₂CO₂H), 5.7 (1H, m, HN-CH), 7.6 (4H, ABq, aromatic C-H), 7.8 (2H, bs, 2xCO₂H), and 8.4 (1H, d, NH-CH).

15 This was dissolved in trifluoroacetic acid (1ml) and stirred at room temperature for 12 hour, and concentrated under reduced pressure to give the desired product (0.15g).

¹H NMR (d₆ DMSO) 3.5 (2H, s, CH₂CO₂H), and 7.6 (4H, AB quartet, aromatic C-H).

Note: NMR in CDCl₃ indicates presence of 5-(4H)-one-2-(4-chlorophenyl)thiazol-4-yl acetic acid tautomer.

20

EXAMPLE 125

25

Preparation of Methyl-2-[5-methyl-2-(4-chlorophenyl)thiazol-4-yl]propionate. (Compound No. 178 in Table I).

A mixture of methyl-(4-bromo-2-methyl-3-oxo) pentanoate (1.0g, 4.5mmol), 4-chlorothiobenzamide (0.7g, 4.5mmol) and ethanol were heated under reflux for 17 hours. The reaction mixture was cooled and
 30 concentrated under reduced pressure. The residue was dissolved in CHCl₃, and washed with NaHCO₃ solution, water, dried (MgSO₄) filtered and concentrated under reduced pressure, to yield the product (0.34g).

¹H NMR (CDCl₃) 1.55 (3H, d, CH₃CH), 2.45 (3H, s, thiazole CH₃), 3.7 (3H, s, CO₂CH₃), 3.95 (1H, q, CH₃CH), and 7.6 (4H, AB quartet, aromatic C-H).

35

EXAMPLE 126

40

Preparation of Methyl-2-[5-bromomethyl-2-(4-chlorophenyl)thiazol-4-yl] propionate. (Compound No. 179 in Table I).

A mixture of methyl-2-(5-methyl-2-(4-chlorophenyl)thiazol-4-yl)propionate (Compound No. 178) (0.5g, 1.7mmol), NBS (0.33g 1.87mmol), benzoyl peroxide (cat.), were irradiated (UV) for 2 hours. The reaction mixture was cooled, filtered, and the filtrate concentrated under reduced pressure. The residue on trituration with hexane gave the product (0.45g).

¹H NMR (CDCl₃) 1.65 (3H, d, CH₃CH), 3.7 (3H, s, CO₂CH₃), 4.0 (1H, q, CH₃CH), 4.8 (2H, AB quartet, CH₂Br), and 7.65 (4H, AB quartet, aromatic C-H).

50

EXAMPLE 127

55

Preparation of compound No. 133 in Table 1.

A mixture of methyl-5-bromo-2-(4-chlorophenyl)thiazol-4-yl acetic ester, (1.0g, 3.0mmol), Lawessons

reagent (2.5g, 6.2mmol) and xylene (10ml) were heated under reflux for 6 hours. The reaction mixture was cooled, filtered and 5g of silica added to the filtrate. The mixture was concentrated under reduced pressure, extracted with CHCl_3 . The combined CHCl_3 extracts were dried, filtered and concentrated under reduced pressure. The product was purified by chromatography (eluted with 50% Et_2O /hexane) to give the desired compound.

^1H NMR (CDCl_3)

4.10 (3H, s, $\text{C}(\text{S})(\text{OCH}_3)$), 4.25 (2H, s, CH_2CO), and 7.6 (4H, AB quartet, aromatic C-H).

Biological Data

The herbicidal activity of the compounds of formula (I) was tested as follows :

The compound in the appropriate concentration was incorporated into a 4% emulsion of methyl cyclohexanone and a 0.4% blend of 3.6 parts Tween 20 and 1 part Span 80. Tween 20 is a Trade Mark for a surface active agent comprising a condensate of 20 molar proportions of ethylene oxide with sorbitan laurate. Span 80 is a Trade Mark for a surface-active agent comprising sorbitan monolaurate. Formulation was effected by dissolving the compound in the requisite amount of solvent/surfactant blend. If necessary glass beads were added, the total liquid volume adjusted to 5ml with water and the mixture shaken to effect complete dissolution of the compound. The formulation so prepared, after removal of beads where necessary, was then diluted to final spray volume (45ml) with water.

The spray compositions so prepared were sprayed onto young pot plants (post-emergence test) at a rate equivalent to 1000 litres per hectare. Damage to plants was assessed 13 days after spraying by comparison with untreated plants, on a scale of 0 to 5 where 0 is 0-10% damage, 1 is 11 to 25% damage, 2 is 26-50% damage, 3 is 51-80% damage, 4 is 81-95% damage and 5 is 96-100% damage.

In a test carried out to detect pre-emergence herbicidal activity, seeds of the test species were placed on the surface of plastic trays of compost and sprayed with the compositions at the rate of 1000 litres per hectare. The seeds were then covered with further compost. 20 Days after spraying, the seedlings in the sprayed plastic trays, are compared to seedlings in untreated control trays, the damage being assessed on same scale of 0-5.

The results of the tests are given in Table III below.

TABLE III

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	TEST PLANTS (SEE TABLE IV)																									
			Sb Rp	Ct	Sy	Mz	Ww	Rc	Bd	Ip	Am	Pi	Ca	Ga	Xa	Xs	Ab	Co	Av	Dg	Al	St	Ec	Sh	Ag	Cn		
4	4	Pre	5	5	4	5	3	4	3	4	2	5	5	5	5	5	-	5	4	1	4	3	3	3	3	5	4	
		Post	2	3	0	4	2	0	0	3	1	4	3	4	3	-	4	3	4	0	1	1	1	0	2	1	3	
17	4	Pre	4	4	3	5	1	3	3	3	3	3	3	4	5	4	-	5	4	1	5	3	2	3	3	4	4	
		Post	0	2	3	3	1	2	0	2	3	3	3	3	4	-	3	3	3	2	2	1	1	3	2	3	3	
19	4	Pre	4	5	1	5	3	4	3	4	3	5	5	5	5	4	-	5	5	0	4	4	2	3	4	5	5	
		Post	2	4	4	4	1	0	2	3	3	4	4	5	4	-	4	4	4	2	3	2	3	3	3	0	4	
20	4	Pre	4	4	1	5	2	4	-	3	4	5	5	5	-	4	-	5	5	1	4	4	2	4	4	5	5	
		Post	3	3	4	4	1	0	0	2	0	5	4	4	4	-	4	4	4	0	0	0	0	1	0	0	0	

TABLE III (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	Sb	Rp	Ct	Sy	Mz	Wz	Rc	Bd	Ip	Am	Pi	Ca	Ga	Xa	Xs	Ab	Co	Av	Dg	Al	St	Ec	Sh	Ag	Cn
24	5	Pre	3	4	0	4	-	5	5	-	0	3	-	2	-	0	-	0	-	-	5	-	5	5	-	5	4
		Post	3	4	1	4	1	3	0	-	2	4	3	4	-	0	-	4	-	1	4	-	5	5	-	3	3
57	4	Pre	2	5	2	4	0	3	4	3	2	-	5	5	-	2	-	5	5	1	3	4	2	4	3	4	4
		Post	1	2	2	4	0	0	1	1	0	4	2	4	4	-	4	2	4	0	0	0	0	0	0	1	0
61	4	Pre	0	2	2	5	0	1	2	4	0	5	5	4	4	4	-	2	5	0	-	2	2	3	3	5	3
		Post	0	1	3	4	1	1	0	1	0	3	3	1	0	-	3	3	3	1	2	2	2	3	3	1	3
63	4	Pre	2	3	1	5	2	3	4	3	3	5	3	3	4	1	-	1	3	1	-	3	3	2	3	3	1
		Post	0	1	0	2	0	0	0	1	3	1	2	2	-	-	1	1	3	-	0	0	0	0	2	0	0
183	4	Pre	4	4	4	4	0	0	0	2	2	4	4	4	4	3	-	3	5	0	0	0	0	0	0	0	3

TABLE III (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	Sb Rp Ct Sy Mz Ww Rc Bd Ip Am Pi Ca Ga Xa Xs Ab Co Av Dg Al St Ec Sh Ag Cn																									
186	4	Pre	5	5	4	5	0	0	0	4	0	4	4	5	5	3	-	5	5	0	1	3	0	0	0	2	4	
		Post	0	0	0	4	0	0	0	1	0	2	0	1	0	-	0	1	0	0	0	0	1	1	0	0	0	
187	4	Pre	0	3	4	4	0	0	2	0	0	4	3	3	-	0	-	1	4	0	0	0	0	0	0	0	3	
193	4	Pre	5	0	2	0	0	2	1	5	0	-	2	5	-	0	-	0	0	3	1	3	0	0	0	-	-	
194	4	Pre	4	4	4	3	0	0	2	2	1	-	2	5	-	0	-	4	4	0	1	0	0	1	0	2	-	
195	4	Post	1	2	2	3	2	0	1	1	3	2	2	2	3	-	0	4	2	1	2	0	2	3	4	1	1	

TABLE III (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	Sb	Rp	Ct	Sy	Mz	Ww	Rc	Bd	Ip	Am	Pi	Ca	Ga	Xa	Ab	Co	Av	Dg	Al	St	Ec	Sh	Ag	Cn	
68	4	Pre	0	0	0	1	1	1	2	0	0	1	3	3	4	0	-	0	1	1	-	2	1	2	3	1	-
		Post	1	1	0	3	0	0	0	0	1	0	1	0	1	-	-	2	0	3	0	0	0	0	3	1	0
70	4	Pre	0	4	2	5	1	2	2	3	0	3	0	0	4	1	-	2	2	0	2	1	2	1	2	-	3
		Post	3	4	3	3	4	1	1	3	0	3	2	-	3	-	3	2	4	2	2	1	0	2	2	2	0
71	4	Pre	3	4	3	5	4	4	5	3	0	3	3	5	4	3	-	5	4	1	4	2	2	3	4	-	5
		Post	2	4	2	4	2	1	0	3	2	4	3	3	4	-	3	3	4	0	2	0	0	1	1	1	2
72	4	Pre	0	4	1	5	3	3	4	3	2	3	4	5	5	3	-	5	5	0	4	2	2	1	3	-	5
		Post	1	3	1	3	3	0	0	3	0	3	3	4	3	-	3	3	3	1	3	0	0	0	0	0	1

TABLE III (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	Sb	Rp	Ct	Sy	Mz	Ww	Rc	Bd	Ip	Am	Pi	Ca	Ga	Xa	Xs	Ab	Co	Av	Dg	Al	St	Ec	Sh	Ag	Cn
76	4	Pre	0	0	0	1	0	0	5	0	0	0	0	-	0	-	-	0	-	0	2	0	0	0	0	-	-
178	3.75	Pre	0	0	0	4	0	0	0	0	2	3	1	0	4	-	1	5	0	0	0	0	0	2	-	-	
		Post	0	0	1	4	0	0	0	0	0	3	0	0	-	3	0	2	0	0	0	0	0	0	0	2	
123	4	Pre	1	2	2	5	0	0	3	1	0	4	4	3	4	0	-	2	0	0	0	0	0	2	-	3	
		Post	0	3	1	4	0	0	0	0	3	1	2	2	-	4	0	4	0	0	0	0	0	0	0	0	
122	3	Pre	1	3	1	5	0	3	3	2	0	4	4	4	4	3	-	2	2	0	0	0	2	3	-	4	
		Post	0	3	4	4	0	0	0	4	0	4	0	2	2	-	4	3	4	0	0	0	0	0	0	1	

TABLE IV

5

Abbreviations used for Test Plants

- 10 Sb - Sugar Beet
 Rp - Rape
 Ct - Cotton
 Sy - Soybean
 Mz - Maize
- 15 Ww - Winter wheat
 Rc - Rice
 Bd - Bidens pilosa
 Ip - Ipomoea purpurea
 Am - Amaranthus retroflexus
- 20 Pi - Polygonum aviculare
 Ca - Chenopodium album
 Ga - Galium aparine
 Xa - Xanthium spinosum
 Xs - Xanthium strumarium
- 25 Ab - Abutilon theophrasti
 Co - Cassia obtusifolia
 Av - Avena fatua
 Dg - Digitaria sanguinalis
 Al - Alopercurus myosuroides
- 30 St - Setaria viridis
 Ec - Echinochloa crus-galli
 Sh - Sorghum halepense
 Ag - Agropyron repens
 Cn - Cyperus rotundus

35 The herbicidal activity of some of the compounds was tested by an alternative method as follows:-Each compound in the appropriate concentration was incorporated into a 4% emulsion of methylcyclohexanone and 0.4% blend of 3.6 parts Tween 20 and 1 part Span 80. Tween 20 is a Trade Mark for a surface active agent comprising a condensate of 20 molar proportions of ethylene oxide with sorbitan laurate. Span 80 is a Trade Mark for a surface-active agent comprising sorbitan monolaurate. Formulation was effected by

40 dissolving the compound in the requisite amount of solvent/surfactant blend. If necessary, glass beads were added, the total liquid volume adjusted to 5ml with water, and the mixture shaken to effect complete dissolution of the compound. The formulation so prepared, after removal of beads where necessary, was then diluted to final spray volume (45ml) with water.

The spray compositions so prepared were sprayed onto young pot plants (post-emergence test) at a

45 rate equivalent to 1000 litres per hectare. Damage to plants was assessed 13 days after spraying by comparison with untreated plants, on a scale of 0 to 9 where 0 is 0% damage, 1 is 1-5% damage, 2 is 6-15% damage, 3 is 16-25% damage, 4 is 26-35% damage, 5 is 36-59% damage, 6 is 60-69% damage, 7 is 70-79% damage, 8 is 80-89% damage and 9 is 90-100% damage.

In a test carried out to detect pre-emergence herbicidal activity, crop seeds were sown at 2 cm depth

50 (i.e. Sb, Ct, Rp, Ww, Mz, Rc, Sy) and weed seeds at 1cm depth beneath compost and sprayed with the compositions at the rate of 1000 litres per hectare. 20 days after spraying, the seedlings in the sprayed plastic trays were compared with the seedlings in unsprayed control trays, the damage being assessed on the same scale of 0 - 9.

The results of the tests are given in Table V below.

55

TABLE V

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	SPECIES (SEE TABLE VI)																									
			Sb	Rp	Ot	Sy	Mz	Rc	Wz	Pi	Ca	Am	Bd	Eh	Ip	Ab	Xa	Xs	Av	Al	Ag	Sh	St	Dg	Ec	Ce		
81	4	Pre	1	1	0	0	2	0	-	3	-	0	0	0	0	0	-	-	0	0	-	0	0	2	0	0		
		Post	5	7	7	9	4	0	0	8	6	7	8	7	4	7	7	-	8	0	0	2	2	4	2	2	0	
90	4	Post	5	5	6	9	3	0	0	7	8	8	7	6	3	2	7	-	7	0	0	5	4	0	0	0	0	
		Pre	1	1	0	1	0	0	0	-	0	-	0	0	0	0	0	-	-	0	0	-	1	0	0	0	1	
94	4	Post	5	6	8	8	6	0	0	3	6	8	8	8	5	2	6	-	7	0	0	3	2	0	3	2	0	
		Pre	0	0	0	2	0	0	0	-	0	-	0	0	0	0	0	-	-	0	0	-	0	0	0	0	0	
71	4	Pre	0	0	0	2	0	0	0	-	0	-	0	0	0	0	0	-	-	0	0	-	0	0	0	0	0	
		Post	5	8	9	9	6	0	0	7	8	8	9	8	4	6	7	-	7	0	0	4	5	5	4	0	0	

TABLE V (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	SPECIES (SEE TABLE VI) Sb Rp Ct Sy Mz Rc Ww Pi Ca Ga Am Bd Eh Ip Ab Xa Xa Av Al Ag Sh St Dg Ec Ce
107	4	Post	5 8 8 8 8 0 3 7 9 8 9 6 5 5 7 - 8 0 0 5 4 2 8 5 0
100	4	Post	7 6 8 8 6 0 0 7 8 8 8 7 4 5 7 - 8 0 0 6 4 3 3 0 0
114	4	Post	5 8 8 8 6 1 2 8 8 8 9 7 5 8 7 - 8 2 0 5 4 4 6 6 0
115	4	Post	5 9 9 9 3 0 2 7 8 8 9 7 5 5 7 - 7 0 0 2 5 2 6 5 0
109	4	Post	6 7 8 9 3 0 0 5 6 8 6 8 4 3 6 - 8 0 2 0 4 4 0 0 0

TABLE V (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	SPECIES (SEE TABLE VI)																								
			Sb	Rp	Ct	Sy	Mz	Rc	Wv	Pi	Ca	Ga	Am	Bd	Eh	Ip	Ab	Xa	Xs	Av	Al	Ag	Sh	St	Dg	Ec	Ce
124	2.2	Pre	0	8	8	9	1	2	7	6	9	9	9	9	4	0	6	9	-	0	3	-	5	0	-	5	5
		Post	1	-	3	7	0	1	2	7	7	-	9	8	3	1	0	-	9	0	0	4	0	0	1	0	0
127	4	Pre	0	2	0	9	0	5	0	2	5	5	0	0	0	0	9	0	-	0	0	-	0	0	0	0	0
		Post	0	0	0	2	0	0	5	0	5	0	5	0	0	0	3	-	3	0	0	0	0	0	0	0	0
128	4	Pre	5	7	0	9	2	3	5	4	7	9	0	5	3	0	4	0	-	0	0	-	0	0	0	0	3
		Post	0	0	4	3	0	0	0	0	2	3	9	1	3	0	3	-	5	0	0	0	3	2	2	1	0
132	1	Pre	0	7	5	9	0	0	5	2	8	7	0	9	0	0	0	9	-	0	0	-	5	0	5	8	2

TABLE V (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	SPECIES (SEE TABLE VI)																									
			Sb	Rp	Ct	Sy	Mz	Rc	Wv	Pi	Ca	Ga	Am	Bd	Eh	Ip	Ab	Xa	Xs	Av	Al	Ag	Sh	St	Dg	Ec	Ce	
147	2.4	Post	6	8	6	9	3	0	4	6	7	7	7	8	7	6	7	-	8	2	2	3	0	4	3	4	0	
148	2.4	Post	6	6	7	8	4	3	4	7	7	6	6	7	6	7	7	-	7	2	2	2	2	4	5	5	2	
149	2.4	Post	5	5	7	8	5	2	5	6	7	7	7	8	6	7	8	-	8	3	2	3	2	2	5	4	0	
150	2.4	Post	5	4	4	8	4	2	3	3	5	5	4	6	3	5	6	-	6	2	3	3	2	2	4	3	0	
162	4	Pre	0	9	9	0	1	0	-	5	-	0	0	0	0	2	0	-	1	0	-	0	0	2	0	0	0	
		Post	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	1	0	0	0	-	0	0	0	0	

TABLE V (Cont/d)

COMPOUND NO.	RATE OF APPLICATION kg/ha	PRE- OR POST EMERGENCE APPLICATION	SPECIES (SEE TABLE VI)																									
			Sb	Rp	Ct	Sy	Mz	Rc	Ww	Pi	Ca	Ga	Am	Bd	Eh	Ip	Ab	Xa	Xs	Av	Al	Ag	Sh	St	Dg	Ec	Ce	
163	4	Pre	0	2	0	9	0	5	3	-	9	-	0	2	0	0	5	0	-	0	5	-	0	0	0	0	3	
		Post	0	0	0	2	0	0	0	0	0	2	2	0	0	0	-	1	0	0	0	-	0	0	0	0		
164	4	Post	2	6	0	0	2	0	0	0	2	4	2	0	0	0	0	-	0	2	0	0	0	0	0	6	0	
168	4	Post	2	0	4	9	0	0	0	2	3	-	3	4	2	3	7	-	-	0	0	0	0	0	0	0		

TABLE VI


5

Abbreviations used for Test Plants in Table V

- 10 Sb - Sugar beet
 Rp - Rape
 Ct - Cotton
 Sy - Soybean
 Mz - Maize
- 15 Rc - Rice
 Ww - Winter wheat
 Bd - Bidens pilosa
 Ip - Ipomoea lacunosa (pre-emergence)
Ipomoea hederacea (post emergence)
- 20 Am - Amaranthus retroflexus
 Pi - Polygonum aviculare
 Ca - Chenopodium album
 Ga - Galium aparine
 Xa - Xanthium spinosum
- 25 Xs - Xanthium strumarium
 Ab - Abutilon theophrasti
 Eh - Euphorbia heterophylla
 Av - Avena fatua
 Dg - Digitaria sanguinalis
- 30 Al - Alopecurus myosuroides
 St - Setaria viridis
 Ec - Echinochloa crus-galli
 Sh - Sorghum halepense
 Ag - Agropyron repens
- 35 Ce - Cyperus esculentes

Claims

- 40 1. A method of killing or controlling unwanted plants by applying to the plant on to a locus thereof an effective amount of a compound of formula (I)



- 55 or a salt thereof: wherein R¹, R², R³, R⁴ and R⁵ are independently selected from hydrogen, hydroxy, alkyl, alkoxy, alkylcarbonyl, halogen, nitrile, nitro, haloalkyl and haloalkoxy;
 R⁶ is hydrogen, hydroxy, amino, lower alkyl, halogen, nitrile, haloalkyl, nitro, aryl, or a group C(O)_mR¹⁰.

wherein R^{10} is hydrogen, alkyl, alkenyl, alkynyl or phenyl any of which may be optionally substituted and m is 1 or 2;

R^7 is hydrogen, lower alkyl, haloalkyl, $C(O)_m R^{10}$ or halogen; or R^6 and R^7 together form an optionally substituted alkylene chain of 2 or 3 carbon atoms; R^8 is hydrogen, lower alkyl or halogen or the group R^7 and R^8 together form an oxo group; and the group CZR^9 is carboxy or an ester thereof, or R^9 is a group SR^{10} or $NR^{11}R^{12}$ wherein R^{11} is hydrogen or alkyl and R^{12} is hydrogen, optionally substituted alkyl, $S(O)_n R^{10}$ wherein n is 0, 1 or 2, or R^9 is a group $-NR^{11}NR^{13}R^{14}$ wherein R^{11} , R^{13} and R^{14} are independently selected from hydrogen or alkyl; or R^9 is a group $-NR^{11}NR^{13}R^{14}R^{20}X^\ominus$ wherein R^{11} , R^{13} , R^{14} and R^{20} are each hydrogen or alkyl, and X^\ominus is an agriculturally acceptable anion; and Z is oxygen or sulphur.

2. A process for selectively controlling the growth of weeds in rice which process comprises applying to the rice or to the growth medium of the rice a compound of formula (I) as defined in claim 1 in an amount sufficient to severely damage or kill weeds but insufficient to damage the rice substantially.

3. A herbicidal composition comprising a compound of formula (I) as defined in claim 1 in combination with an agriculturally acceptable carrier or excipient.


4. A composition according to claim 3 wherein the group CZR^9 is carboxy or an ester thereof.

5. A composition according to claim 3 or claim 4 wherein R^6 is lower alkyl or halogen.

6. A composition according to claim 5 where R^6 is chlorine or bromine.


7. A composition according to any one of claims 3 to 6 wherein R^9 is other than hydrogen.

8. A compound of formula (II)



wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 and R^9 are as defined in claim 1; and R^{17} is halogen nitrile, nitro, hydroxy, amino or mono or dialkyl amino.

9. A process for preparing a compound of formula (II) as defined in claim 8 which process comprises either (a) where R^{17} is halogen reacting a compound of formula (III)




wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , and R^8 are as defined in claim 1 with chlorine or bromine. and thereafter if desired converting the group $CZOH$ to a different group CZR^9 as defined in claim 1 or

(b) where R^{17} is nitrile, subjecting a compound of formula (I) wherein R^6 is alkoxycarbonyl to conventional procedures for transforming an alkoxycarbonyl group to a $-CN$ group.

(c) where R^{17} is nitro, amino or mono or dialkyl amino reacting an ester derivative of a compound of formula (III) with a mild nitrating agent and thereafter if desired carrying out one or more of the following steps:


- (i) hydrogenating the nitro group R^{17} to an amino group;
- (ii) alkylating the amino group R^{17} ; and
- (iii) converting the ester group to a different groups CZR^9 .

10. A compound of formula (V):




15 wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^8 , and R^9 and Z are as defined in claim 1.

11. A process for preparing a compound of formula (V) as defined in claim 10 which process comprises reacting a compound of formula (VI):



30 wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined in claim 1; with a compound of formula (VII):



40 wherein R^8 , R^9 and Z are as defined in claim 1 and X^1 is a leaving group.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 31 1472

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
D,A	GB-A-1 022 750 (SHELL INTERNATIONALE RESEARCH MAATSCHAPPIJ N.V.) * Claims * ---	1-11	A 01 N 43/78 C 07 D 277/20 C 07 D 277/60
A	US-A-3 749 787 (WALTER HEPWORTH et al.) * Column 1, lines 41-64; column 2, lines 54,56; claims 1-2 * & GB-A-1 099 389 (Cat. D) ---	1-7	
A	GB-A-1 147 626 (JOHN WYETH & BROTHERS LTD) * Claims 1-3 * ---	1-7	
A	GB-A-1 137 529 (JOHN WYETH & BROTHERS LTD) * Claims 1-2,9 * ---	1-7	
A	CHEMICAL ABSTRACTS, vol. 77, no. 19, 6th November 1972, page 21, abstract no. 122118e, Columbus, Ohio, US; F.J. McEVOY et al.: "2-Phenyl-4H-cyclopentathiazole-4-carboxylic acids as potential antiinflammatory agents", & J. MED. CHEM. 1972, 15(8), 850-1 * Abstract * -----	10-11	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A 01 N C 07 D
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 02-02-1990	Examiner RAVANEL C.M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	